

Complete nucleotide sequence of *Saccharomyces cerevisiae* chromosome X

F.Galibert^{1,2}, D.Alexandraki³, A.Baur⁴, E.Boles⁴, N.Chalwatzis⁴, J.-C.Chuat¹, F.Coster⁵, C.Cziepluch⁶, M.De Haan⁷, H.Domdey⁸, P.Durand⁹, K.D.Entian¹⁰, M.Gatius¹, A.Goffeau⁵, L.A.Grivell⁷, A.Hennemann¹⁰, C.J.Herbert¹¹, K.Heumann¹², F.Hilger⁹, C.P.Hollenberg¹³, M.-E.Huang¹, C.Jacq¹⁴, J.-C.Jauniaux⁶, C.Katsoulou³, L.Kirchrath¹³, K.Kleine¹², E.Kordes⁶, P.Kötter¹⁰, S.Liebl¹², E.J.Louis¹⁵, V.Manus¹, H.W.Mewes¹², T.Miosga⁴, B.Obermaier^{8,16}, J.Perea¹⁴, T.Pohl¹⁷, D.Portetelle⁹, A.Pujol⁶, B.Purnelle⁵, M.Ramezani Rad¹³, S.W.Rasmussen¹⁸, M.Rose¹⁰, R.Rossau¹⁹, I.Schaaff-Gerstenschläger⁴, P.H.M.Smits⁷, T.Scarcez¹⁹, N.Soriano¹, D.Tovan¹⁴, M.Tzermia³, A.Van Broekhoven¹⁹, M.Vandenbol⁹, H.Wedler²⁰, D.Von Wettstein¹⁸, R.Wambutt²⁰, M.Zagulski^{11,21}, A.Zöllner¹² and L.Karpfinger-Hartl¹²

¹UPR 41 CNRS Recombinations Génétiques, Faculté de Médecine, 2 avenue du Professeur Léon Bernard, F-35043 Rennes Cedex, France.

²Foundation for Research and Technology Hellas, Institute of Molecular Biology and Biotechnology, PO Box 1527, Heraklion, GR-71110 Crete, Greece. ³Institut für Mikrobiologie und Genetik, Technische Hochschule Darmstadt, Schnittspahnstrasse 10, D-64287 Darmstadt, Germany. ⁴Unité de Biochimie Physiologique, Université Catholique de Louvain, Place Croix du Sud 2, Bâtiment 20, B-1348 Louvain-La-Neuve, Belgium. ⁵Immunovirologie Abteilung 0610 and Virologie Appliquée à l'Oncologie Unité INSERM U375, Deutsches Krebsforschungszentrum, D-69120 Heidelberg, Germany. ⁶University of Amsterdam, Section for Molecular Biology, Kruislaan 318, NL-1098 SM Amsterdam, The Netherlands. ⁷Genzentrum, Institut für Biochemie, Würmlstrasse 221, D-81373 München, Germany. ⁸Unité de Microbiologie, Faculté des Sciences Agronomiques de Gembloux, avenue Maréchal Juin 6, B-5030 Gembloux, Belgium. ⁹Institut für Mikrobiologie, J.W.Goethe-Universität Frankfurt, Marie-Curie-Strasse 9, Geb. N250, D-60439 Frankfurt/Main, Germany. ¹⁰UPR 2420 CNRS Centre de Génétique Moléculaire, Bâtiment 26, Avenue de la Terrasse, F-91198 Gif-sur-Yvette cedex, France. ¹¹MIPS am Max-Planck-Institut für Biochemie, D-82152 Martinsried bei München, Germany. ¹²Institut für Mikrobiologie der Heinrich-Heine-Universität Düsseldorf, Geb. 26.12, Universitätsstrasse 1, D-40225 Düsseldorf, Germany. ¹³URA 1302 CNRS Génétique Moléculaire, Ecole Normale Supérieure, 46 rue d'Ulm, F-75230 Paris Cedex 05, France. ¹⁴Yeast Genetics, Institute of Molecular Medicine, John Radcliffe Hospital, Headington, Oxford OX3 9DU, UK. ¹⁵GATC GmbH, Gesellschaft für Analyse Technik und Consulting, Fritz-Arnold-Strasse 23, D-78467 Konstanz, Germany. ¹⁶Carlsberg Laboratory, Department of Physiology, Gamle Carlsberg vej 10, Valby, DK-2500 Copenhagen, Denmark. ¹⁷Innogenetics, Industriepark Zwijnaarde 7, Box 4, B-9052 Ghent, Belgium and ¹⁸AGON GmbH, Gesellschaft für molekularbiologische Technologie mbH, Glienicke Weg 185, D-12489 Berlin, Germany

¹⁹Present address: MediGene GmbH, Lochhamer Strasse 11, D-82152 Martinsried bei München, Germany

²¹Present address: Institute of Biochemistry and Biophysics, 5a Pawlowskiego St., 02-106 Warsaw, Poland

²Corresponding author

The complete nucleotide sequence of *Saccharomyces cerevisiae* chromosome X (745 442 bp) reveals a total of 379 open reading frames (ORFs), the coding region covering ~75% of the entire sequence. One hundred and eighteen ORFs (31%) correspond to genes previously identified in *S.cerevisiae*. All other ORFs represent novel putative yeast genes, whose function will have to be determined experimentally. However, 57 of the latter subset (another 15% of the total) encode proteins that show significant analogy to proteins of known function from yeast or other organisms. The remaining ORFs, exhibiting no significant similarity to any known sequence, amount to 54% of the total. General features of chromosome X are also reported, with emphasis on the nucleotide frequency distribution in the environment of the ATG and stop codons, the possible coding capacity of at least some of the small ORFs (<100 codons) and the significance of 46 non-canonical or unpaired nucleotides in the stems of some of the 24 tRNA genes recognized on this chromosome.

Keywords: chromosome X/gene duplication/open reading frame/*Saccharomyces cerevisiae*/tRNA

Introduction

The traditional methods of genetic analysis involve tracing modified phenotypes back to genotypic alterations. The limit of this approach is an imperceptible modification of the phenotype. The international yeast genome systematic sequencing programme launched in 1989 by the European Communities, aiming at establishing the complete genetic information of bakers' yeast, *Saccharomyces cerevisiae*, has demonstrated the limitations of classical genetics. The pilot sequencing of chromosome III (Oliver *et al.*, 1992) has demonstrated that disruption of a large number of the newly revealed open reading frames (ORFs) does not result in any phenotypic alteration. Subsequent systematic sequencing of seven more chromosomes (Barrell *et al.*, 1994; Dietrich *et al.*, 1994; Dujon *et al.*, 1994; Feldmann *et al.*, 1994; Johnston *et al.*, 1994; Bussey *et al.*, 1995; Murakami *et al.*, 1995) has confirmed that a large proportion of the novel genes cannot be assigned any known function, while on the other hand a large number of proteins unrelated to database entries are being discovered. Last but not least, it stems from numerous cytological studies of chromosome behaviour during the vegetative and meiotic cell cycle that a chromosome is more than its mere genetic content. By making available the complete

Table 1. Estimated overall accuracy of chromosome X sequence

	Total bp verified	Number of modified nt ^a			Error rate (%)
		M	G	T	
Overlap between regions	46 455	11	13	24	0.52
Resequenced regions ^b	~50 000	10	7	17	0.34

^aM, mismatch; G, gap; T, total mismatches plus gaps.

^bOccasional overlaps between verification clone sequences were excluded from the calculations.

DNA sequence of a chromosome, parameters not entirely confined to its role as carrier of genetic information may be exposed for analysis. A survey of a new object is thus provided, even though all the topological implications of the results cannot be fully grasped at the present stage and must await at least the completion of the yeast genome enterprise. This paper describes the DNA sequence of chromosome X.

Results

Assembly of the sequence

The sequence was determined from a set of 26 partially overlapping cosmids selected on the basis of an *EcoRI* map based on a cosmid contig of chromosome X (Huang *et al.*, 1994a). These cosmids were distributed within a consortium of 15 contractors. The telomeres were independently isolated and sequenced. While the left-telomere-containing clone was found to overlap with the left terminal cosmid of the chromosome, this was not so at the other end, where no overlap was detected between the right-most cosmid and a right-telomere-containing clone 9.0 kb in size. The missing portion (a few kb) was PCR-amplified from a yeast S288C genomic DNA template using primers designed from sequences flanking the gap. When all bases had been determined by each contractor and each sequencing strategy had been approved by the DNA coordinator, ensuring that the sequence had been independently determined on each strand with sufficient overlap between all the subclones, the sequences were considered as final and entered into the MIPS data library for assembly. Partial sequences of chromosome X have been published independently by some of the authors of this work (Huang *et al.*, 1994b, 1995; Miosga *et al.*, 1994a,b,c, 1995; Purnelle *et al.*, 1994; Vandenbol *et al.*, 1994, 1995; Rasmussen, 1995; Zagulski *et al.*, 1995).

Verification of the sequence

Quality controls were performed concomitantly with sequence assembly. The aim of the project was to keep the error rate as low as possible, with a target $<10^{-4}$. Three procedures were employed to track down errors, including checking sequencing strategy by the coordinator, matching overlapping portions sequenced by independent contractors and finally random resequencing (see Materials and methods for details). The results of the last two procedures are shown in Table I. From these data, the error rate of the yeast chromosome X sequence presented in this paper can be estimated to be 0.4%, a value of the same order as that reported in similar studies.

General organization of chromosome X

Analysis of the entire nucleotide sequence of chromosome X (745 442 bp) confirms the general features of chromosome organization observed in other systematically sequenced yeast chromosomes. The coding region occupies 74.04% of the sequence, 36.59% and 37.45% on the Watson and Crick strand, respectively.

The average base composition is 38.9% G+C. As expected, the coding regions have a higher than average G+C content (40.2%) than the non-coding (35.6%). The distribution of dinucleotide frequencies over the whole chromosome is the same in the coding and the non-coding regions of either strand. The deviations of the frequencies of complementary dinucleotide pairs tend to occur in the same direction. In contrast to what was reported for chromosomes XI and II, the homopurine pairs do not seem to be in excess in the coding region of either strand (Figure 1). Some compositional periodicity has been noted, at least in the case of chromosomes XI and II, with waves of G+C-rich regions correlating with waves of high gene density. By using the same algorithm, a similar G+C pattern emerges with chromosome X, especially in the right-hand part of the chromosome. This pattern correlates rather well with the gene density plot, as illustrated by the two deep depressions around 200 kb and 470 kb in Figure 2.

Telomeres and centromere

The telomere regions of chromosome X are similar to the other sequenced yeast telomeres. Adjacent to the C_{1-3} A repeat at the left telomere are a Y' element (coordinates 61–6931) and the core X element (7305–7767) shared by most if not all yeast telomeres (Louis *et al.*, 1994; Pryde *et al.*, 1995). However, the X–Y' junction does not contain the usual subtelomeric repeats STR-D, STR-C, STR-B and STR-A, but instead has (6998–7224) part of a copy (Louis and Haber, 1991) of the fourth intron of cytochrome *b* encoded by mitochondrial DNA (Delehdode *et al.*, 1989). A copy of *bi4* is also found at the left telomere of chromosome IX (Louis and Haber, 1991; Barrell *et al.*, 1994). In fact, the left ends of chromosomes IX and X share a large, nearly identical block of sequence similarity spanning >21 kb. The right telomere of chromosome X is more conventional, with a core X element (744 593–745 052) and the STR-D, STR-C, STR-B and STR-A elements adjacent to the TG_{1-3} repeats (745 357–end). The core X elements of both ends contain the *ARS1* consensus and the Abf1p binding site found in most core Xs. These elements that are shared by most ends may have functional significance. The right telomere region is analogous to several other sequenced telomeres (II right




Fig. 1. Distribution of dinucleotide frequencies in the coding and non-coding regions of the two strands of chromosome X. Vertical bars show relative deviations [i.e. (observed-expected)/expected]. Expected frequencies are calculated from mononucleotide frequencies. Complementary pairs are arranged as mirror images. The four self-complementary pairs are placed in the central part.




Fig. 2. Compositional variation and gene density distribution along chromosome X. Top: compositional variation calculated as in Dujon *et al.* (1994). Each point represents the average G+C composition calculated from the third base of each codon. Bottom: gene density expressed as the fraction of nucleotides within ORFs in sliding windows of 30 kb. The position of the centromere is indicated by an arrow.

and left, V right and left, VI left, VIII right and left, IX right, XI left) over the last 3–4 kb.

The centromere of chromosome X of strain R95-4A, a derivative of S288C, was isolated by Hieter *et al.* (1985) by selection of yeast DNA fragments capable of suppressing lethality of the *SUP11* gene in high copy number. Comparison of this sequence with that reported in the present

paper shows complete identity and enables location of the chromosome X centromere at positions 435 996–436 112. *CEN10* conforms to the consensus structure established for other centromeres.

ORFs and their predicted protein products

By definition, an ORF is considered from its first in-phase ATG codon. Only those ORFs containing at least 99 contiguous sense codons following an ATG, and not entirely contained within a longer ORF in a different reading frame or on the other DNA strand, have been retained for further analysis. The special case of ORFs shorter than 100 codons is described below. A total of 379 ORFs were recorded in the entire chromosome X using this principle (Table II), leaving aside the retroposons, i.e. a density of one ORF/1967 bp. Twelve of these ORFs are interrupted by introns. Table II includes 39 partially overlapping ORFs. Ten are on the same DNA strand, all others being antiparallel overlaps. Informatic and statistical analysis revealed that ORFs both shorter than 150 codons and with a codon adaptation index (CAI) (Sharp and Li, 1987) <0.11 may correspond to randomly occurring ORFs rather than to real genes (Dujon *et al.*, 1994). If these criteria are applied to the ORFs identified in chromosome X, 23 of the 379 ORFs are questionable genes. Thirteen of these belong to the set of partially overlapping ORFs. However, three genes of known function (*YAP17*, *STE18* and *RPL46*) fall into this category as well, making the border between ORF and gene even more elusive. Taking into account the physical position and ATG environment may help tell which ORFs are genes.

Comparison of the nucleotide sequence and of the predicted protein products with public database entries reveals that 118 ORFs (31%) correspond to genes previously identified in *S.cerevisiae*. All other ORFs represent

Table II. List of ORFs longer than 99 sense codons, known genes and other genetic elements of chromosome X

Nomenclature		Size (aa)	Coordinates	Locus	CAI	Fasta score	Description (nature of element, function or similarity of product)/Comment		
Working	Official		1	60					
			61	6931			left telomere sequence (complement TG ₁₋₃)		
			61	6931			Y' element		
J0202	YJL225c	1504	469	6130	0.11		probable nucleotide-binding protein, TMM 1+1 (intron from 4582 to 4969)	E	
			6998	7224			copy of part of b14 intron from cytochrome <i>b</i> gene (mitochondrial DNA)		
			7305	7767			cutc X element		
J0208	YJL223c	120	8779	9138	0.65	534 (5381)	similar to PAG1 protein (PIR: S48516)	B	
J0213	YJL222w	1549	11475	16121	0.16	5326 (7778)	similar to carboxypeptidase Y-sorting protein PEP1 (PIR: S25329), TMM 3+1	B	
J0218	YJL221c	589	16770	18536	0.25	2459 (3094)	similar to α -glucosidase MAL35 (PIR: S46183), TMM 1+0	B	
J0220	YJL220w	150	18243	18692	0.10		hypothetical protein, TMM 2+1	B	
J0222	YJL219w	567	19974	21197	0.17	2913 (2955)	similar to hexose transport protein LGT3 (PIR: S45153), TMM 8+1	E	
J0224	YJL218w	196	21973	22560	0.11	453 (943)	similar to galactoside <i>O</i> -acetyltransferase (SW: P07464), TMM 1+0	C	
J0226	YJL217w	198	23133	23726	0.12		hypothetical protein	F	
J0228	YJL216c	581	24344	26086	0.23	2229 (3095)	similar to α -glucosidase (PIR: S45157), TMM 1+0	C	
J0231	YJL215c	119	26415	26771	0.10		hypothetical protein, ?	F	
J0232	YJL214w	569	26887	28593	0.20	2953 (3021)	probable hexose transport protein HXT6 (PIR: S45159), TMM 11+1	B	
J0234	YJL213w	331	32163	33155	0.14		hypothetical protein	F	
J0236	YJL212c	799	33853	36249	0.18	1610 (4357)	similar to <i>S.pombe</i> ISP4 (PIR: S45161), TMM 10+1	D	
J0238	YJL211c	147	36760	37200	0.10		hypothetical protein, ?	F	
J0240	YJL210w	271	36919	37731	<i>CRT1</i>	0.09	CRT1 protein (PIR: S27422)	A	
J0242	YJL209w	654	38005	39966	<i>CBP1</i>	0.15	CBP1 protein (PIR: S05829)	A	
J0310	YJL208c	329	40197	41183	<i>NUC1</i>	0.14	nuclease NUC1 precursor, mitochondrial (PIR: S05888)	A	
J0312	YJL207c	2014	41392	47433		0.14	hypothetical protein, TMM 4+1	E	
J0316	YJL206c	758	47662	49935		0.15	hypothetical protein, TMM 1+1	E	
J0318	YJL205c	187	50632	51192		0.14	hypothetical protein	F	
J0320	YJL204c	645	51216	53150		0.16	hypothetical protein	F	
J0322	YJL203w	280	53340	54179	<i>SPP91</i>	0.14	pre-mRNA splicing factor SPP91 (PIR: S23553)	A	
J0323	YJL202c	115	53945	54289		0.12	hypothetical protein, TMM 1+1	E	
J0325	YJL201w	599	54378	56174		0.15	hypothetical protein	F	
J0327	YJL200c	789	56446	58812		0.22	2130 (3762)	similar to mitochondrial acinifate hydratase (GB: U17709)	C
J0330			59099	59171			tRNA ^{Thr}		
J0332			59471	59782			δ terminant		
J0334	YJL199c	108	59857	60180		0.09	hypothetical protein, ?	F	
J0336	YJL198w	881	60842	63484	0.18	2799 (4318)	similar to YCR037c (PIR: S46633), TMM 13+1	C	
J0340	YJL197w	1254	63803	67564	0.14	535 (6137)	probable ubiquitin-carboxyl terminal hydrolase (SW: P35123)	D	
J0343	YJL196c	310	67851	68780	0.13	924 (1753)	similar to sterol isomerase SUR4 (PIR: S46638), TMM 5+0	C	
J0345	YJL195c	233	69242	69940		0.11	hypothetical protein, TMM 2+0	F	
J0347	YJL194w	513	69336	70874	<i>CDC6</i>	0.13	cell division control protein CDC6 (PIR: S46640)	A	
J0349	YJL193w	402	71364	72569		0.10	447 (2131)	similar to SLY41 protein (PIR: S46641), TMM 6+1	D
J0351	YJL192c	234	72711	73412		0.16	hypothetical protein, TMM 2+0	E	
J0353	YJL191w	1380	73785	74606	<i>CRY2</i>	0.59	ribosomal protein S14eB (intron from 73795 to 74202) (PIR: S46643)	A	
J0355	YJL190c	130	74911	75300	<i>RPS24</i>	0.81	ribosomal protein S15eE (PIR: A23082)	A	
J0360	YJL189w	51	75931	76469	<i>RPL46</i>	0.92	ribosomal protein L39c (intron from 75937 to 76322) (EMBL: X01963)	B	
J0403	YJL188c	102	76203	76508		0.15	hypothetical protein	F	
J0406	YJL187c	819	76804	79260	<i>SWE1</i>	0.13	protein kinase SWE1 (PIR: S40400), TMM 1+0	A	
J0409	YJL186w	586	80152	81909		0.16	1039 (3004)	similar to TTP1 protein (PIR: S45870), TMM 2+0	C
J0415	YJL185c	293	82095	82973		0.11	hypothetical protein	F	
J0420	YJL184w	123	83445	83813		0.08	hypothetical protein, ?	F	
J0425	YJL183w	422	84065	85330		0.18	hypothetical protein, TMM 1+0	E	
J0430	YJL182c	105	85435	85749		0.08	hypothetical protein, TMM 1+0, ?	E	
J0435	YJL181w	611	85657	87489		0.11	443 (2950)	hypothetical protein, similar to J1575, TMM 1+1	F
J0486	YJL180c	325	87583	88557	<i>ATP12</i>	0.12	ATP12 protein precursor (PIR: A39736)	A	
J0488	YJL179w	109	88784	89110		0.15	hypothetical protein	F	
J0490	YJL178c	196	89282	89869		0.17	hypothetical protein, TMM 1+0	E	
J0493	YJL177w	184	90782	91651		0.68	825 (827)	ribosomal protein L17c (intron from 91091 to 91407) (PIR: S38012)	B
J0495	YJL176c	825	92052	94526	<i>SWI3</i>	0.15	transcription factor SWI3 (PIR: S26706)	A	
J0502	YJL175w	170	94045	94554		0.12	hypothetical protein, TMM 3+0	E	
J0504	YJL174w	276	95088	95915	<i>KRE9</i>	0.16	secretory pathway protein KRE9 precursor (PIR: S23891), TMM 1+0	A	
J0506	YJL173c	122	96160	96525	<i>REB3</i>	0.14	replication factor A chain 3 (PIR: C37281)	A	
J0510	YJL172w	411	97729	99456	<i>CPST</i>		Gly-X carboxypeptidase precursor (PIR: S16693)	A	
J0512	YJL171c	396	99699	100886		0.22	478 (1923)	hypothetical protein, similar to YBR162C (PIR: S46033), TMM 2+0	D
J0514	YJL170c	183	101145	101693		0.13	hypothetical protein, TMM 2+0	E	
J0517	YJL169w	122	102090	102455		0.15	hypothetical protein, TMM 2+0	E	
J0520	YJL168c	733	102221	104419		0.14	258 (3593)	similar to human A11.1 zinc finger motif (PIR: A44264)	D
J0525	YJL167w	282	105005	106060	<i>FPP1</i>		farnesyl-pyrophosphate synthetase (SW: A34441), TMM 1+1	A	
J0526	YJL166w	94	106425	106706		0.21	QCR8	ubiquitin-cytochrome <i>c</i> reductase subunit VIII (PIR: S48138)	A
J0531	YJL165c	855	106888	109452	<i>HAL5</i>	0.13	HAL5 protein (PIR: S48240)	A	
J0541	YJL164c	397	109960	111150	<i>SRA3</i>	0.18	protein kinase, cAMP-dependent, catalytic chain 1 (PIR: A27070)	A	
J0544	YJL163c	555	111662	113326		0.08	hypothetical protein, TMM 11+1	E	
J0549	YJL162c	482	114177	115622		0.14	hypothetical protein	F	

Table II. Continued

Nomenclature	Size (aa)	Coordinates	Locus	CAI	FastA score	Description (nature of element, function or similarity of product)/Comment		
Working Official								
JH55H		115932	116003			tRNA ^{Glu}		
JH552	YJL161w	180	117238	117777	0.19	hypothetical protein, TMM 1+1	E	
JH555	YJL160e	180	118281	118819	0.15	326 (751)	similar to PIR1 protein (chr XI) (PIR: S33650)	C
JH558	YJL159w	310	120443	121372	0.47	577 (1162)	similar to PIR2 protein (chr XI) (PIR: S33651)	C
JH561	YJL158e	227	121964	123644	0.59	521 (976)	similar to PIR2 protein (chr XI) (PIR: S33651)	C
JH565	YJL157e	830	123535	126124	FAR1	factin arrest protein FAR1 (SW: S13341)	A	
JH57H	YJL156r	687	126589	128649	0.13		hypothetical protein, TMM 1+1	E
JH575	YJL155e	452	128985	130341	FBP26	fructose-2,6-bisphosphate 2-phosphatase (PIR: A42569)	A	
JH58H	YJL154c	944	130801	133632	VPS35	vacuolar protein-sorting protein VPS35 (PIR: S31293)	A	
JH61D	YJL153e	555	134032	135696	INO1	myo-inositol-1-phosphatase (PIR: A30902), TMM 2+1	A	
JH628	YJL152w	119	135871	136227	0.07		hypothetical protein, TMM 1+H, ?	E
JH63H	YJL151e	133	136372	136470	0.16		hypothetical protein, TMM 2+H	E
JH632	YJL150w	100	136820	137119	0.09		hypothetical protein, TMM 1+H, ?	E
JH634	YJL149w	663	137076	139064	0.16	296 (3276)	hypothetical protein, similar to YD93D2.D6c (IGB: S51858), TMM 1+0	E
JH635		139458	139647	SNR190		SnR 190 small nuclear RNA		
JH636		139263	140301	SNR128		SnR 128 small nuclear RNA		
JH637	YJL148w	233	140134	140832	0.20		hypothetical protein	F
JH639	YJL147e	382	141119	142264	0.13		hypothetical protein	F
JH642	YJL146w	469	142989	144395	0.11		hypothetical protein, TMM 1+H	E
JH644	YJL145w	294	144857	145738	0.22		hypothetical protein	F
JH646	YJL144w	104	146156	146367	0.07		hypothetical protein, ?	F
JH648	YJL143w	158	146798	147271	MIM17	mitochondrial inner membrane protein MIM17 (PIR: S46257), TMM 1+1	A	
JH650	YJL142r	130	147519	147908	0.06		hypothetical protein, TMM 3+1, ?	E
JH652	YJL141r	817	147667	150087	YAK1	protein kinase YAK1 (PIR: A32582), TMM 1+H	A	
JH654	YJL140w	221	150658	151320	RPB4	DNA-directed RNA polymerase II chain RPB4 (PIR: A32491)	A	
JH657	YJL139e	428	151413	152696	YUR1	YUR1 protein (PIR: S26856), TMM 1+0	A	
JH66H	YJL138c	395	153204	154388	TIF2	translation initiation factor eIF-4A(IGB: X12814)	A	
JH663	YJL137r	381	154685	155824	0.14	445 (1978)	hypothetical protein, similar to YKR058w (PIR: S38134)	D
JH664	YJL136c	87	156247	156970	0.60		ribosomal protein S21e (from 156487 to 156946)	B
JH666	YJL135w	105	157574	157888	0.14		hypothetical protein	F
JH671	YJL134w	409	157885	159111	0.11	1298 (2332)	hypothetical protein, similar to YKR053c (PIR: S38127), TMM 4+1	E
JH675	YJL133w	314	160316	161257	MRS3	splicing protein MRS3, mitochondrial (PIR: S01267)	A	
JH678	YJL132w	751	161611	163861	0.12		hypothetical protein, TMM 1+1	E
JH682	YJL131e	356	163978	165045	0.12		hypothetical protein	F
JH686	YJL130c	2214	165423	172064	URA2	pyrimidine synthesis protein URA2 (PIR: S05767), TMM 1+1	A	
JH689	YJL130Ac	115	171926	172929	0.06		hypothetical protein, (intron from 172082 to 172740), ?	F
JH693	YJL129r	1235	173299	177003	TRK1	potassium transport protein, high-affinity (PIR: S05849), TMM 8+1	A	
JH699	YJL128r	668	177797	179801	PBS2	polymyxin B resistance protein kinase (PIR: A32714)	A	
JH702	YJL127e	640	181999	183918	SPT10	regulatory protein SPT10 (PIR: S47865)	A	
JH706	YJL126w	307	184199	185119	0.12	309 (1519)	hypothetical protein, similar to L9638.5 (IGB: U0902)	F
JH71H	YJL125c	383	185229	186377	0.14		hypothetical protein	F
JH714	YJL124c	172	186828	187343	0.16		hypothetical protein	F
JH718	YJL123r	478	187706	189139	0.15		hypothetical protein	F
JH723	YJL122w	175	189415	189939	0.21		hypothetical protein	F
JH731	YJL121c	238	190076	190789	RPE1	ribose-5-phosphate 3-pyrimidase (IGB: 83571)	A	
JH734	YJL120w	0.17	190721	191041	0.14		hypothetical protein, TMM 1+1	E
JH738	YJL119c	0.17	191274	191594	0.13		hypothetical protein, TMM 1+H	E
JH742	YJL118w	219	191338	191994	0.09		hypothetical protein, TMM 1+1	E
JH744	YJL117w	311	192230	193162	0.19		hypothetical protein, TMM 2+H	E
JH748	YJL116c	337	193562	194572	0.25	1091 (1566)	hypothetical protein, similar to YKR042w (PIR: S38114), TMM 1+0	E
JH755	YJL115w	279	195985	196821	ASF1	ASF1 protein (PIR: S07664), TMM 1+1	A	
JH76D		197011	197083			tRNA ^{Asp}		
JH765		197193	197242			delta remnant		
JH77H		197243	197613			subunit LTR of Ty4		
JH775		414	197613	198854	Ty4A_JL	Ty4A_JL protein		
JH78D		1803	197613	203022	Ty4B_JL	Ty4B_JL protein		
JH785		203098	203468			subunit LTR of Ty4		
JH790		203503	203814			delta remnant		
JH795		203815	204092			delta remnant		
JH799		204431	204502			tRNA ^{Asp}		
JH802	YJL112w	714	205001	207142	0.12	229 (4303)	probable G-protein, beta-transducin type (PIR: B48088)	D
JH804	YJL111w	550	207573	209222	0.19	1754 (2527)	probable rho-terminin of the TCC-1 ring complex, similar to mouse CCT7 (PIR: S43058)	C
JH806	YJL110c	551	209621	211273	GZF3	GATA zinc finger protein 3 (IGB: X86353)	E	
JH808	YJL109r	1769	211699	217005	0.17		hypothetical protein, TMM 5+1	B
JH811	YJL108c	383	217404	218552	0.17		hypothetical protein, TMM 8+1	E
JH813	YJL107c	387	218552	219712	0.13		hypothetical protein	F
JH817	YJL106w	645	221086	223020	SME1	probable protein kinase SME1 (PIR: S20138), TMM 1+H	A	
JH819	YJL105w	560	224751	226430	0.03	586 (2734)	hypothetical protein, similar to YKR029c (PIR: S38101), TMM 1+0	E
JH822	YJL104w	149	227023	227469	0.09		hypothetical protein, ?	F

Table II. Continued

Nomenclature	Size (aa)	Coordinates	Locus	CAI	Fasta score	Description (nature of element, function or similarity of product)/Comment	
J0823		228122 228297	<i>SNR37</i>			SnR 37 small nuclear RNA	
J0824	YJL103c	618 228724 230577		0.12	253 (12980)	probable haem dependent regulatory protein, similar to S46116	D
J0826	YJL102w	819 230997 233453	<i>MEF2</i>	0.13		translation elongation factor G homologue, MEF2, mitochondrial IPIR: S43748). TMM 1+1	A
J0829		233635 233707				tRNA ^{Asp}	
J0832	YJL101c	678 234019 236052	<i>GSH1</i>	0.14		glutamate-cysteine ligase (PIR: S28648). TMM 2+1	A
J0834	YJL100w	607 236959 238779		0.11		hypothetical protein	F
J0838	YJL099w	746 239110 241347	<i>CSD3</i>	0.12		CSD3 protein (GB: U15603)	A
J0840	YJL098w	1058 241778 244951		0.15	1625 (4985)	hypothetical protein, similar to YKR028w (GB: X85H21)	F
J0902	YJL097w	217 245287 245937		0.18		hypothetical protein, TMM 6+0	E
J0904	YJL096w	224 245997 246668		0.13		hypothetical protein, TMM 2+0	E
J0906	YJL095w	1478 246950 251383	<i>BCK1</i>	0.12		protein kinase BCK1 IPIR: S201171	A
J0909	YJL094c	873 251519 254137		0.13	264 (14290)	probable transport protein, similar to PIR: A42111. TMM 13+1	E
J0911	YJL093c	691 254435 256507	<i>TOK1</i>	0.12		TOK1, outwardly rectifying potassium channel protein, TMM 10+0 F	F
J0913	YJL092w	1174 257118 260639	<i>RADH1</i>	0.13		helicase RADH (PIR: S46586)	A
J0916	YJL091c	498 260778 262271		0.13		hypothetical protein, TMM 8+1	E
J0918	YJL090c	764 262455 264746		0.14		hypothetical protein	F
J0922	YJL089w	829 265621 268107	<i>SIP4</i>	0.14		SIP4 protein, probable regulatory protein (GB: U17643). TMM 2+1	A
J0924	YJL088w	440 268188 269507	<i>ARG3</i>	0.16		ornithine carbamyltransferase IPIR: S10058). TMM 1+1	A
J0927	YJL087c	827 269700 272180	<i>TRL1</i>	0.16		tRNA ligase (PIR: A29917). TMM 1+1	A
J0930	YJL086c	122 272176 272541		0.11		hypothetical protein, TMM 1+0	E
J0932	YJL085w	623 272522 274390		0.16		hypothetical protein	F
J0934	YJL084c	1046 274560 277697		0.13	1555 (4683)	hypothetical protein, similar to YKR021W (PIR: S38090)	F
J1002	YJL083w	604 278536 280347		0.09	596 (2822)	hypothetical protein, similar to YKR019c (PIR: S38088)	F
J1007	YJL082w	731 280880 283072		0.17	2652 (3586)	hypothetical protein, similar to YKR018c (PIR: S38087). TMM 1+1	E
J1012	YJL081c	489 283500 284966	<i>ACT3</i>	0.13		actin-related protein IPIR: S47608	A
J1017	YJL080c	1222 285256 288921	<i>SCP160</i>	0.33		SCP160 protein, histone-like protein (PIR: S37492)	A
J1022	YJL079c	299 289573 290469		0.30	670 (1268)	hypothetical protein, similar to YKR013W (PIR: S38082). TMM 1+0	C
J1027	YJL078c	881 291034 293676		0.15	597 (3322)	hypothetical protein, similar to YKR013W (PIR: S38082). TMM 2+1	D
J1033	YJL077c	131 294364 294756		0.08		hypothetical protein, TMM 1+1, ?	E
J1038	YJL076w	1189 294940 298506		0.15	345 (4906)	putative protein-binding protein, similar to YKR010c (PIR: S25814)	D
J1044	YJL075c	138 298158 298571		0.11		hypothetical protein, TMM 1+0	F
J1049	YJL074c	1230 298855 302544		0.18	605 (5561)	probable purine nucleotide-binding protein, similar to SMC1 (PIR: S41804). TMM 1+0 D	D
J1083	YJL073w	692 302735 304810		0.14		hypothetical protein, TMM 1+1	E
J1086	YJL072c	213 304919 305557		0.12		hypothetical protein, TMM 1+0	E
J1091	YJL071w	574 305827 307548		0.12	314 (2803)	similar to acetyl-glutamate synthase (GB: L35484). TMM 1+1	D
J1095	YJL070c	888 307669 310332		0.14	441 (4614)	hypothetical protein, similar to YBR284w (PIR: S47120). TMM 1+1	E
J1098	YJL069c	594 310620 312401		0.17		hypothetical protein	F
J1102	YJL068c	299 312714 313610		0.20	525 (1572)	similar to human esterase D (SW: P10768)	D
J1107	YJL067w	116 313779 314126		0.12		hypothetical protein, TMM 1+1	E
J1111	YJL066c	252 313812 314567		0.16		hypothetical protein	F
J1115	YJL065c	167 314752 315252		0.11		hypothetical protein	F
J1120	YJL064w	131 314870 315262		0.12		hypothetical protein, TMM 1+1	E
J1125	YJL063c	238 315457 316170	<i>MRPL8</i>	0.09		ribosomal protein L17, mitochondrial (PIR: S47128)	A
J1132	YJL062w	830 316979 319468		0.12		hypothetical protein, TMM 9+1	E
J1135	YJL061w	713 319711 321849		0.16		hypothetical protein	F
J1138	YJL060w	444 323081 324412		0.21	662 (2193)	probable amino acid transferase, similar to (PIR: S52790)	D
J1139	YJL059w	408 324659 325882		0.12		hypothetical protein, TMM 6+1	E
J1141	YJL058c	543 325940 327568		0.12	1119 (12465)	purine nucleotide binding protein, similar to YBR270c (PIR: S46151). TMM 1+0	C
J1143	YJL057c	667 327816 329816		0.14		hypothetical protein, TMM 1+1	E
J1145	YJL056c	880 330129 332768		0.16	436 (4257)	probable regulatory protein, similar to mouse Kr2 protein (PIR: S10549), leucine zipper D	D
J1148	YJL055w	245 333052 333786		0.14		hypothetical protein	F
J1150	YJL054w	478 333960 335393		0.15		hypothetical protein	F
J1152	YJL053w	379 335593 336729	<i>PEP8</i>	0.14		PEP8 protein (PIR: S48882)	A
J1154	YJL052w	332 337966 338961	<i>TDH1</i>	0.86		glyceraldehyde-3-phosphate dehydrogenase 3 (PIR: A10372). TMM 1+1	A
J1156	YJL051w	822 339482 341947		0.12		hypothetical protein, TMM 3+0	E
J1158	YJL050w	1073 342217 345435		0.20	971 (15214)	viral mRNA translation inhibitors SK12 (GB: D29641)	D
J1162	YJL049w	450 345668 347017		0.16		hypothetical protein	F
J1164	YJL048c	396 347145 348332		0.14	344 (11921)	hypothetical protein, similar to YBR273c (PIR: S46154)	F
J1166	YJL047c	842 349278 351803		0.12		hypothetical protein	F
J1171	YJL046w	451 351955 353307		0.12	302 (12257)	similar to lipotein-protein ligase <i>A. Ecofi</i> IPIR: A54035)	D
J1173		353939 354027				tRNA ^{Val} (small intron)	
J1177		354233 354555				su10 δ	
J1179		354539 354870				su10 δ	
J1185		355069 355140				tRNA ^{Asp}	
J1190		355151 355222				tRNA ^{Asp}	
J1194	YJL045w	634 355719 357620		0.16	2721 (13048)	similar to succinate dehydrogenase flavoprotein IPIR: S34793)	B
J1202	YJL044c	458 357998 359371	<i>GYP6</i>	0.16		GTPase-activating protein GYP6 IPIR: S31061). TMM 1+0	A

Table II. Continued

Nomenclature		Size	Coordinates	Locus	CAI	FastA score	Description (nature of element, function or similarity of product)/Comment	
Working	Official	(aa)						
J1204	YJL043w	257	359825 360595		0.09		hypothetical protein	F
J1206	YJL042w	1398	360944 365137	<i>MIP1</i>	0.15		microtubule-associated protein (GB: X84652)	B
J1207	YJL041w	823	365479 368065	<i>NSP1</i>	0.16		nucleoskeletal-like protein NSP1 (PIR: S14055) (intron from 365480 to 365597)	B
J1216	YJL039c	1683	368446 373494		0.15		hypothetical protein, TMM 4+1	E
J1221			374119 374190				tRNA ^{Asp}	
J1226			374201 374272				tRNA ^{Asp}	
J1230			374539 374630				soln δ	
J1232	YJL038c	219	374813 375469		0.10	405 (1049)	similar to J1234, TMM 3+0	E
J1234	YJL037w	224	376357 377028		0.11	405 (1049)	similar to J1232, TMM 2+1	E
J1240			378055 378128				tRNA ^{Glu}	
J1244	YJL036w	423	378520 379788		0.15		hypothetical protein	F
J1246	YJL035c	250	379947 380696		0.12		hypothetical protein	F
J1248	YJL034w	682	381022 383067	<i>KAR2</i>	0.44		nuclear fusion protein KAR2 precursor (PIR: A32366), TMM 1+1	A
J1250	YJL033w	770	383532 385841		0.20	530 (3629)	similar to <i>E. coli</i> SrmB RNA helicase (SW: P21507)	D
J1252	YJL032w	104	386043 386354		0.15		hypothetical protein	F
J1254	YJL031c	390	386066 386935	<i>BET4</i>	0.15		geranylgeranyl transferase α chain (PIR: S48301)	A
J1256	YJL030w	196	387352 387939	<i>MAD2</i>	0.12		MAD2 protein (PIR: S48302)	A
J1258	YJL029c	822	388083 390548		0.13	317 (4044)	similar to <i>C. elegans</i> T05G5.8 protein (PIR: S41008)	F
J1263			390738 390810				tRNA ^{Met}	
J1267	YJL028w	111	391006 391338		0.07		hypothetical protein, TMM 2+0, ?	E
J1269	YJL027c	138	391531 391944		0.08		hypothetical protein, ?	F
J1271	YJL026w	399	392099 393295	<i>RNR2</i>	0.50		ribonucleoside-diphosphate reductase small chain (PIR: A26916), TMM 1+1	A
J1273	YJL025w	514	393662 395203	<i>RRN7</i>	0.13		RRN7 protein (PIR: S50785)	A
J1274	YJL024c	194	395623 396287		0.14	229 (920)	related to mouse clathrin associated protein 19 (intron from 396189 to 396265) (PIR: A40535)	D
J1278			396421 396491				tRNA ^{Gly}	
J1282	YJL023c	347	397053 398093		0.13		hypothetical protein	F
J1284	YJL022w	402	397804 398109		0.10		hypothetical protein, TMM 1+1, ?	F
J1286	YJL021c	365	398635 399729		0.13		hypothetical protein	F
J1305	YJL020c	771	399789 402101		0.14	206 (3404)	glutamic acid rich protein precursor (<i>Plasmodium falciparum</i>) (PIR: A54514)	D
J1310	YJL019w	620	402588 404447		0.12		hypothetical protein, TMM 1+0	E
J1315	YJL018w	104	404321 404632		0.16		hypothetical protein	F
J1320	YJL017w	325	405278 406252		0.13		hypothetical protein	F
J1326	YJL016w	171	406447 406959		0.16		hypothetical protein	F
J1331	YJL015c	124	406834 407205		0.12		hypothetical protein	F
J1336	YJL014w	534	407246 408847	<i>BIN2</i>	0.23		chaperonin of the TCP-1 ring complex, TMM 1+1, similar to mouse CCT3 (PIR: S43062)	B
J1341	YJL013c	515	409184 410728		0.13	475 (2454)	similar to protein kinase B1B1 (Yeast chr 7) (GB: I.M32027)	D
J1345	YJL012c	648	411143 413086		0.25		hypothetical protein	F
J1349	YJL011c	161	413975 414457		0.12		hypothetical protein	F
J1352			414653 414725				tRNA ^{Leu}	
J1355			415618 415724				tRNA ^{Trp} (small intron)	
J1357	YJL010c	666	417252 419249		0.17		hypothetical protein	F
J1369	YJL009w	108	419542 419865		0.16		hypothetical protein, TMM 1+1	F
J1374	YJL008c	568	419647 421350		0.20	1219 (2622)	probable chaperonin of the TCP-1 ring complex, similar to mouse CCT8 (PIR: S52867)	C
J1379	YJL007c	104	422388 422699		0.13		hypothetical protein, TMM 1+0	E
J1385			422624 422696				tRNA ^{Met}	
J1390	YJL006c	323	422828 423796		0.11		hypothetical protein, TMM 1+0	E
J1395			424119 424202				tRNA ^{Leu}	
J1401	YJL005w	2026	424844 430921	<i>CYR1</i>	0.12		adenylate cyclase (PIR: A24776)	A
J1402	YJL004c	203	431279 431887		0.09		hypothetical protein, TMM 4+0	E
J1403	YJL003w	118	432331 432684		0.10		hypothetical protein, TMM 1+0, ?	E
J1404	YJL002c	476	432911 434338	<i>OST1</i>	0.16		α subunit, oligosaccharyltransferase (GB: Z46719), TMM 2+0	A
J1407	YJL001w	193	435032 435610	<i>PRE3</i>	0.17		multicatalytic endopeptidase complex chain PRE3 (PIR: S43669), TMM 1+0	A
			435996 436018	<i>CDEIII</i>			centromere	
			436022 436104	<i>CDEII</i>			centromere	
			436105 436112	<i>CDEI</i>			centromere	
J1409	YJR0011w	602	436489 438294		0.12	257 (2951)	similar to <i>C. elegans</i> , hypothetical protein (PIR: S42372), TMM 10+1	E
J1411	YJR002w	593	438551 440329		0.17		hypothetical protein	F
J1415	YJR003c	539	440683 442399		0.13		hypothetical protein	F
J1418	YJR004c	650	442598 444547	<i>AGAL1</i>	0.13		α -agglutinin (PIR: S22835), TMM 2+0	A
J1422	YJR005w	445609	447708	<i>YAP80</i>			clathrin-associated protein complex β chain homolog (PIR: S12934), TMM 1+1	A
J1427	YJR006w	487	448888 450348		0.16		hypothetical protein	F
J1429	YJR007w	304	450706 451617	<i>SGJ2</i>	0.37		translation initiation factor eIF-2 α chain (PIR: A32108)	A
J1431	YJR008w	338	452116 453129		0.14		hypothetical protein	F
J1433	YJR009c	332	453372 454367	<i>TDH2</i>	0.90		glyceraldehyde-3-phosphate dehydrogenase (PIR: S40915)	A
J1436	YJR010w	511	455925 457457	<i>MET3</i>	0.29		sulfate adenylyltransferase (PIR: S00906)	A
J1438	YJR011c	261	458330 459112		0.14		hypothetical protein	F
J1440	YJR012c	207	459484 460104		0.12		hypothetical protein, TMM 1+0	E

Table II. Continued

Nomenclature	Size (aa)	Coordinates	Locus	CAI	Fasta score	Description (nature of element, function or similarity of product)/Comment	
Working	Official						
J1444	YJRH13w	305	460363-461277		D.11	hypothetical protein, TMM 5+1	E
J1446	YJRH14w	198	461516-462109		D.22	hypothetical protein	F
J1448	YJRH15w	510	462408-463937		D.13 1380 [2637]	similar to SNG1 gene (yeast chr 7) (GB: X7492H), TMM 5+1	C
J1450	YJRH16c	585	464141-465895	<i>ILV3</i>	H.38	dihydroxy-acid dehydratase (PIR: S43744)	A
J1452	YJRH17c	190	466211-466780	<i>ESS1</i>	H.12	ESS1 protein (PIR: S07867)	A
J1454	YJRH18w	120	466473-466832		H.08	hypothetical protein, TMM 1+1, ?	E
J1456	YJRH19c	349	466922-467968		D.11 222 [1776]	similar to <i>E.coli</i> acyl-CoA thioesterase	D
J1458	YJRH20w	111	467688-468017		H.11	hypothetical protein, TMM 1+1	E
J1462	YJRH21c	292	468310-469266	<i>MER2</i>	H.11	nicotinic recombination protein MER2 (intron from 468871 to 468950) (PIR: A40271)	A
J1464	YJRH22w	128	469414-469797		H.13	hypothetical protein	F
J1470	YJRH23c	133	469494-469892		H.09	hypothetical transport protein, TMM 2+1, ?	E
J1545	YJRH24c	244	469920-470651		D.12	hypothetical protein	F
J1550	YJRH25c	177	470828-471358		G.17 313 [922]	similar to human 3-hydroxyanthranilate 3,4-dioxygenase (PIR: A5407H)	D
J1553			472150-472487			δ , LTR of Ty1	
J1555		440	472447-473766		H.14 1990 [2005]	TyA protein	
J1560		174	472447-477712		H.15 824 [8276]	TyB protein	
J1563			477738-478071			δ , LTR of Ty1	
J1565		440	478031-479350		H.15 1991 [1997]	TyA protein	
J1570		174	478031-483296		D.14 825 [8277]	TyB protein	
J1573			483322-483659			δ , LTR of Ty1	
J1575	YJRH30c	745	483649-485883		H.11 443 [3553]	hypothetical protein, similar to J0435	F
J1580	YJRH31c	1408	486276-490499		H.13 317 [6683]	hypothetical protein, similar to YEJ102w (PIR: S24168), TMM 6+1	E
J1585	YJRH32w	393	490768-491946		H.19 468 [1962]	hypothetical protein, similar to L8167.24 (PIR: S48567)	F
J1590	YJRH33c	1357	492068-496138		H.14 310 [6771]	hypothetical protein	F
J1604	YJRH34w	008	496370-496693	<i>PET191</i>	H.12	PET191 protein (PIR: S28924)	A
J1606	YJRH35w	1085	497042-500296	<i>RAD26</i>	H.13	probable helicase RAD26 (SW: P40352), TMM 1+1	A
J1608	YJRH36c	892	500403-503078		H.11	hypothetical protein, TMM 1+1	E
J1610	YJRH37w	127	502789-503169		O.11	hypothetical protein	F
J1612	YJRH38c	120	503400-503759		H.09	hypothetical protein, TMM 2+0, ?	F
J1614	YJRH39w	1121	503623-506985		D.13	hypothetical protein, TMM 2+1	E
J1616	YJRH40w	779	507433-509769		D.14 788 [3956]	similar to mouse chloride channel protein (GB: D17521), TMM 7+1	D
J1622	YJRH41c	1174	509929-513450		D.14	hypothetical protein, TMM 2+1	E
J1624	YJRH42w	744	513742-515973		H.13	hypothetical protein, TMM 1+0	E
J1626	YJRH43c	350	516151-517200		H.14	hypothetical protein	F
J1631			517500-517571			tRNA ^{Met}	
J1634			517645-517786			δ remnant	
J1637	YJRH44c	140	518453-518872		H.15	hypothetical protein, TMM 4+0	E
J1639	YJRH45c	654	519328-521289	<i>SSC1</i>	D.52	heat shock protein 70-related protein SSC1 precursor, mitochondrial (PIR: A32493)	A
J1641	YJRH46w	604	521735-523546		H.11	hypothetical protein, TMM 1+1	E
J1647			523699-523780			tRNA ^{Ser}	
J1651	YJRH47c	157	524598-525068	<i>ANB1</i>	H.00	translation initiation factor eIF-5A.2 (PIR: B40259)	A
J1653	YJRH48w	109	526022-526348	<i>CYC1</i>	H.37	cytochrome c isoform 1	A
J1655	YJRH49c	530	526574-528163	<i>UTR1</i>	H.13	UTR1 protein (PIR: S46589), TMM 1+1	A
J1657	YJRH50w	235	528384-529088	<i>UTR3</i>	H.00	UTR3 protein (PIR: S46590)	A
J1659	YJRH51w	501	529548-531050	<i>OSM1</i>	H.17	OSM1 protein precursor (PIR: S46591), TMM 1+0	A
J1661			531202-531361			δ remnant	
J1663			531515-531585			tRNA ^{Gly}	
J1665	YJRH52w	565	531749-533443	<i>RAD7</i>	H.14	RAD7 protein (PIR: A25226)	A
J1667	YJRH53w	574	533714-535435		H.15	hypothetical protein	F
J1669	YJRH54w	497	535743-537233		H.13 725 [2484]	hypothetical protein, similar to YM9827H5c (GB: Z47816), TMM 4+0	E
J1670			538242-538313			tRNA ^{Arg}	
J1705	YJRH55w	164	538459-538950	<i>HIT1</i>	D.13	HIT1 protein (PIR: S30869)	A
J1706a			540453-540783			salto δ	
J1706b			540786-541114			salto δ	
J1707			541195-541266			tRNA ^{Asp}	
J1710	YJRH56c	236	541482-542289		G.10	hypothetical protein	F
J1713			542643-542731			tRNA ^{Pro} (small intron)	
J1715	YJRH57w	216	543749-544396	<i>CDC8</i>	G.15	dTMP kinase (PIR: A00683)	A
J1720	YJRH58c	147	544422-544862	<i>YAP17</i>	G.08	clathrin-associated protein 17 (PIR: C40535)	A
J1725	YJRH59w	818	545474-547927		H.16 1251 [3786]	similar to serine/threonine specific protein kinase (PIR: S30135), TMM 1+0	D
J1730	YJRH60w	351	548446-549408	<i>CBF1</i>	H.14	centromere-binding protein CBF1 (PIR: A36300)	A
J1736	YJRH61w	935	550198-553002		D.13	hypothetical protein, TMM 1+1	E
J1742	YJRH62c	457	553166-554536	<i>NTA1</i>	H.12	amino-terminal amidase NTA1 (PIR: S47938)	A
J1747	YJRH63w	125	554882-555256	<i>RPA12</i>	H.00	DNA-directed RNA polymerase I chain A1.2 (PIR: A48007), TMM 1+0	A
J1752	YJRH64w	562	555600-557286		H.22 1704 [2637]	probable chaperonin of the TCP-1 ring complex, similar to mouse CCT5 (PIR: S43060), TMM 1+0	C
J1760	YJRH65c	449	557499-558845		H.20 1499 [2153]	similar to actin-like protein Act 2 (Hissin yeast) (PIR: A41790), TMM 1+1	C
J1803	YJRH66w	2470	559003-566512	<i>TOR1</i>	D.14	TOR1 protein (PIR: S43940), TMM 3+1	A
J1805	YJRH67c	141	566709-567031		H.14	hypothetical protein	F

Table II. Continued

Nomenclature	Size (aa)	Coordinates	Locus	CAI	Fasia score	Description (nature of element, function or similarity of product)/Comment		
Working Official								
J1808	YJR068w	353	567330	568388	<i>RFC2</i>	0.18	replication factor C chain RFC2 (PIR: S45531)	A
J1811	YJR069c	197	568496	569086		0.20	hypothetical protein	F
J1814	YJR070c	325	569311	570285		0.40	hypothetical protein	F
J1818	YJR071w	122	570092	570457		0.10	hypothetical protein. ?	F
J1821	YJR072c	385	570657	571811		0.17	847 (1816) similar to <i>C.elegans</i> protein C34E10 (GB: U10402)	F
J1824	YJR073c	206	572005	572622	<i>PEM2</i>	0.17	methylene-fatty-acyl-phospholipid synthase (PIR: B28443), TMM 3+1	A
J1827	YJR074w	218	572782	573435		0.15	hypothetical protein	F
J1830	YJR075w	396	573668	574855		0.18	209 (2020) similar to mannosyltransferase (PIR: S22701), TMM 2+0	D
J1833	YJR076c	415	575044	576288	<i>CDC11</i>	0.17	cell division control protein CDC11 (PIR: S40911)	A
J1837	YJR077c	311	576945	577877	<i>MIR1</i>	0.36	phosphate transport protein, mitochondrial (PIR: S12318), TMM 1+1	A
J1840	YJR078w	453	578547	579905		0.13	514 (2251) similar to mouse indoleamine 2,3-dioxygenase (PIR: JH0492)	D
J1843	YJR080w	109	579892	580923		0.11	hypothetical protein (intron from 580035 to 580739), TMM 1+0	E
J1847	YJR081c	394	580122	581303		0.14	hypothetical protein	F
J1854	YJR082c	113	581604	581942		0.15	hypothetical protein	F
J1857	YJR083c	309	582298	583224		0.11	hypothetical protein	F
J1860	YJR084w	423	583420	584688		0.10	hypothetical protein	F
J1863	YJR085c	105	584810	585124		0.14	hypothetical protein, TMM 2+0	E
J1866	YJR086w	110	585755	586084	<i>STE18</i>	0.10	STE18 protein (PIR: B30102)	A
J1870	YJR087w	116	586087	586431		0.10	hypothetical protein, TMM 2+0. ?	E
J1875	YJR088c	292	586185	587060		0.17	hypothetical protein	F
J1880	YJR089w	954	587405	590266		0.13	hypothetical protein	F
J1885	YJR090c	1151	590562	594014	<i>GRR1</i>	0.12	GRR1 protein (PIR: A41529), TMM 1+1	A
J1890	YJR091c	1091	594751	598023		0.15	593 (4842) hypothetical protein, similar to YP9499.01c (PIR: S54067)	F
J1901	YJR091Ac	200	597437	598035		0.15	ATP/GTG binding site motif A	E
J1905	YJR092w	1320	598809	602768		0.15	hypothetical protein	F
J1911	YJR093c	327	602916	603896	<i>FIP1</i>	0.12	component of pre-mRNA polyadenylation factor	B
J1916	YJR094c	360	604265	605344	<i>IME1</i>	0.18	meiosis-inducing protein IME1 (PIR: S31137)	A
J1921	YJR095w	322	609466	610431	<i>ACR1</i>	0.21	ACR1 protein (PIR: S43280), TMM 2+1	A
J1926	YJR096w	282	610888	611733		0.22	431 (1491) probable reductase protein, similar to GB: A32950	D
J1931	YJR097w	172	612106	612621		0.13	hypothetical protein	F
J1936	YJR098c	655	612882	614846		0.15	hypothetical protein	F
J1941	YJR099w	236	615266	615973	<i>YUHI</i>	0.11	ubiquitin carboxyl-terminal hydrolase YUHI (GB: S51332), TMM 1+0	A
J1946	YJR100c	327	616044	617024		0.10	hypothetical protein	F
J1950			617609	617709			IRNA ^{low} (small intron)	
J1952	YJR101w	266	617924	618721		0.11	hypothetical protein	F
J1957	YJR102c	202	618850	619455		0.13	hypothetical protein	F
J1962	YJR103w	564	620444	622135	<i>URA8</i>	0.16	C'TP synthase URA8 (PIR: S42580), TMM 2+0	A
J1968	YJR104c	154	622242	622703	<i>SOD1</i>	0.38	superoxide dismutase (Cu-Zn) (PIR: A36171)	A
J1973	YJR105w	340	623270	624289		0.37	hypothetical protein	F
J1978	YJR106w	725	624527	626701		0.10	hypothetical protein	F
J1983	YJR107w	328	627030	628013		0.13	hypothetical protein, TMM 12+1	E
J1988	YJR108w	123	628403	628771		0.14	hypothetical protein	F
J2002	YJR109c	1118	629279	632632	<i>CPA2</i>	0.24	large subunit of arginine specific carbamoyl-phosphate synthase (PIR: A01199)	A
J2007	YJR110w	688	633306	635369	<i>CPA1</i>	0.16	small subunit of arginine specific carbamoyl-phosphate synthase (PIR: B33478)	A
J2009	YJR111c	283	635549	636397		0.12	hypothetical protein	F
J2011	YJR112w	201	636721	637323		0.09	hypothetical protein	F
J2020	YJR113c	247	637926	638666		0.10	204 (1185) similar to ribosomal protein S7 (<i>Bacillus stearothermophilus</i>) (PIR: JG0008)	D
J2024	YJR114w	130	638350	638739		0.11	hypothetical protein, TMM 1+0	F
J2027	YJR115w	169	639633	640139		0.10	hypothetical protein	E
J2031	YJR116w	279	640516	641352		0.14	hypothetical protein, TMM 2+1	E
J2032	YJR117w	453	641698	643056		0.27	hypothetical protein, TMM 5+1	E
J2033	YJR118c	203	643184	643792		0.19	hypothetical protein, TMM 3+1	E
J2035	YJR119c	728	644998	646181		0.15	776 (3828) similar to human retinoblastoma binding protein 2 (GB: S66431)	D
J2039	YJR120w	116	646817	647164		0.07	hypothetical protein. ?	F
J2041	YJR121w	511	647298	648830	<i>ATP2</i>	0.42	H ⁺ -transporting ATP synthase β chain precursor (PIR: S27278)	A
J2043	YJR122w	497	649467	650957		0.15	hypothetical protein	F
J2045	YJR123w	125	651592	652266	<i>RPS5</i>	0.75	ribosomal protein S5	A
J2046	YJR124c	448	652586	653929		0.14	hypothetical protein, TMM 9+1	E
J2048	YJR125c	408	654431	655654		0.17	283 (1775) hypothetical protein, similar to L8167.6 yeast protein (PIR: S48557)	F
J2050	YJR126c	811	655948	658388		0.13	521 (3981) similar to human prostate-specific membrane antigen (SW: Q04609), TMM 1+0	D
J2052	YJR127c	1380	658611	662750	<i>ZMS1</i>	0.12	ZMS1 protein (PIR: S43751), TMM 4+1	A
J2059	YJR128w	119	662612	662968		0.06	hypothetical protein. ?	F
J2060			663440	663633	<i>SNR3</i>		SnR 3 small nuclear RNA	
J2061	YJR129c	339	663694	664710		0.11	hypothetical protein, TMM 1+0	E
J2063	YJR130c	639	664912	666828		0.13	1778 (3174) similar to TUB1 3' region (GB: S49644)	C
J2110	YJR131w	549	667335	668981	<i>MNS1</i>	0.14	α -mannosidase MNS1 (PIR: A39345), TMM 1+0	A
J2112	YJR132w	1048	669213	672356		0.15	hypothetical protein, TMM 2+1	E
J2118	YJR133w	209	672682	673308		0.28	hypothetical protein	F
J2120	YJR134c	707	673423	675543		0.15	343 (3092) similar to human TATA element modulatory factor (PIR: A47212)	D

Table II. *Continued*

Nomenclature	Size (aa)	Coordinates	Locus	CAI	FastA score	Description (nature of element, function or similarity of product)/Comment		
Working Official								
J2122	YJR135c	239	675753 676469		0.12	hypothetical protein	F	
J2124	YJR136c	421	677135 678397		0.10	hypothetical protein	F	
J2126	YJR137c	1442	678651 682976	0.25	1054 (6897)	similar to ferredoxin sulfate reductase (SW: P30008)	D	
J2129	YJR138w	1584	684258 689009		0.14	hypothetical protein	F	
J2132	YJR139c	359	689139 690215	<i>HOM6</i>	0.47	homoserine dehydratase (PIR: S33317). TMM 1+1	A	
J2161	YJR140c	1648	690444 695387		0.14	hypothetical protein, TMM 1+1	E	
J2166	YJR141w	347	695597 696637		0.13	hypothetical protein, TMM 1+1	E	
J2171	YJR142w	342	696832 697857		0.15	hypothetical protein	F	
J2176	YJR143r	762	698020 700305	<i>PMT4</i>	0.22	PMT4 protein (PIR: S51284). TMM 8+1	A	
J2181	YJR144w	269	700573 701379	<i>MGM101</i>	0.16	MGM101 protein (PIR: S34849)	A	
J2186	YJR145r	261	7011721 702759	<i>RPS7A</i>	0.69	ribosomal protein S4ec10 lintron form 702490 to 7027451 (PIR: S20054)	A	
J2200	YJR146w	117	703576 703926		0.07	hypothetical protein, ?	F	
J2204	YJR147w	358	703887 704960		0.12	235 (1782)	similar to heat shock transcription factor 8 (PIR: S25481)	D
J2209	YJR148w	376	705435 706562		0.19	1584 (1900)	similar to TWT1 yeast protein (PIR: S48989)	C
J2213	YJR149w	404	706851 708062		0.14	462 (1937)	similar to 2-nitropropane dioxygenase (PIR: S50891)	D
J2217	YJR150c	298	708505 709398		0.30		hypothetical protein, TMM 2+0	E
J2223	YJR151r	1161	711949 715431		0.23	614 (14382)	similar to human mucin (PIR: A40963). TMM 2+0	D
J2230	YJR152w	543	719357 720985	<i>DAL5</i>	0.16		allantoate permease (PIR: A28671). TMM 6+1	A
J2235	YJR153w	361	722506 723588		0.17	907 (11643)	similar to polygalacturonase (PIR: S28771). TMM 1+0	C
J2240	YJR154w	346	725475 726512		0.13		hypothetical protein	F
J2245	YJR155w	288	727036 727959		0.15	1334 (1439)	similar to yeast aryl-alkanol dehydrogenase (PIR: S51335)	B
J2250	YJR156r	340	728268 729287		0.53	1784 (1790)	similar to thiamine-repressed amt-1 protein (PIR: S48548). TMM 1+0	B
J2255	YJR157w	120	730206 730565		0.13		hypothetical protein, TMM 1+0	F
J2260	YJR158w	567	732131 733831		0.16	1893 (13036)	similar to hexose transport protein HXT7 (PIR: S43186). TMM 9+1	C
J2395	YJR159w	357	735735 736805	<i>SOR1</i>	0.22		sorbitol dehydrogenase (IGB: L11139)	B
J2400	YJR160c	602	737702 739507		0.13	2585 (14048)	similar to sugar transport protein ISW: P38156). TMM 7+1	C
J2403	YJR161c	383	742542 743690		0.14	1845 (12635)	similar to YB8L (SW: P38363). TMM 3+1	E
			744593 745052				ene X element	
			745053 745356				STR-D, C, B and A elements	
J2420	YJR162c	116	744605 744952		0.14	422 (804)	similar to YKW5 (SW: P36030)	F
			745357 745442				right telomere sequence	

Last column: status of the protein deduced from each putative gene. The categories A (fully known) to F (unknown) are defined in the text. The self FastA score of the predicted protein is in parentheses. An accession number in one of the public databases [PIR, Swiss-Prot (SW), GenBank (GB) and EMBL] is indicated. Abbreviations: TMM: Transmembrane motif, integral+ peripheral; ?: questionable gene. ORF YJL093c is categorized as F, as it was discovered and sequenced during the systematic sequencing of chromosome X and found to correspond to no known gene. It was subsequently biologically characterized as a potassium channel (Keichum *et al.*, 1995).

novel putative yeast genes whose function will have to be determined experimentally. However, 57 of these (another 15% of total) encode proteins that show significant similarity to a protein of known function from yeast or other organisms, thus providing some indication as to their function. The 204 (54%) remaining ORFs exhibit no significant similarity to known sequences (FastA score <200). Motif searches have shown that 91 of the latter have some particular protein signature, mostly a structure suggestive of transmembrane domains (Table II).

An approximately equal number of ORFs is observed on each DNA strand. The mean ORF size is 482 codons (1446 bp), the longest (YJR066w) reaching 2470 codons. The mean size of inter-ORF regions, disregarding one in each pair of overlapping ORFs, is 602 bp for terminator-promoter combinations (WW and CC in Figure 3). For divergent promoters (DP) and convergent terminators (CT), the mean size is 725 bp and 311 bp, respectively. This striking difference in inter-ORF size between divergent promoters versus convergent terminators may be indicative of more important sequence requirements in promoter regions for the regulation of gene expression. An exception is the contiguity of the two ORFs YJL108c and YJL107c. The TGA stop codon of the latter overlaps the ATG of the former, so that both codons share TG. This peculiarity was carefully checked by oligo-primed sequencing in

either direction on cosmid DNA. The two ORFs in their integrity are translated from a single transcript of ~3 kb (Rasmussen, 1995).

Environment of ATG and stop codons

Compilation of a large number of sequence data surrounding the initiation codon AUG has revealed that these sequences are not random and that higher eukaryotes have in common the consensus sequence GCC(A/G)CCATGG (Kozak, 1987). In the case of the budding yeast, another consensus (A/Y)A(A/Y)A(A/Y)AATGGTCT has been proposed (Hinnebusch and Liebman, 1991).

We examined the 318 chromosome X ORFs longer than 150 codons, in all probability corresponding to real genes, to test this consensus. Table III shows the frequency of the different nucleotides, as determined by tabulating positions -8 to +7 relative to ATG. A χ^2 test was performed at each position to test the non-randomness of this distribution, taking into account the G+C content of the chromosome. At all positions except -5 the distribution was found to be non-random. As these calculations are based on all the ORFs of a chromosome, regardless of their expression level, rather than on a selected subset, the following consensus sequence might be more appropriate: AAANAAAATGGCTG. The chances of a random distribution at each position is <5%, or even 1%




Fig. 3. Mean size and distribution of inter-ORF regions of chromosome X. WW: promoter/terminator combination on Watson strand; CC: promoter/terminator combination on Crick strand; DP: divergent promoters; CT: convergent terminators. The numbers indicate on top line the mean size, on bottom line the distribution of each configuration.

Table III. Initiation and stop codon environment

ATG environment													
	-8	-7	-6	-5	-4	-3	-2	-1	ATG	+4	+5	+6	+7
A	0.396	0.393	0.368	0.349	0.399	0.569	0.403	0.456	ATG	0.318	0.283	0.324	0.327
G	0.164	0.160	0.211	0.135	0.148	0.195	0.119	0.145	ATG	0.296	0.129	0.151	0.299
C	0.173	0.192	0.176	0.220	0.189	0.113	0.252	0.173	ATG	0.132	0.362	0.182	0.129
T	0.267	0.255	0.245	0.296	0.264	0.123	0.226	0.223	ATG	0.254	0.343	0.343	0.242
χ^2	7.978	9.616	10.015	7.370	10.060	104.811	30.284	27.741	ATG	20.165	61.227	8.750	22.695

TAG stop codon environment												
	-5	-4	-3	-2	-1	TAG	+4	+5	+6	+7	+8	+9
A	0.380	0.268	0.310	0.394	0.296	TAG	0.408	0.282	0.380	0.437	0.366	0.282
G	0.127	0.183	0.253	0.211	0.211	TAG	0.211	0.127	0.293	0.211	0.197	0.141
C	0.183	0.197	0.169	0.085	0.113	TAG	0.113	0.197	0.183	0.056	0.169	0.239
T	0.310	0.352	0.268	0.310	0.380	TAG	0.268	0.394	0.197	0.296	0.268	0.338
χ^2	2.975	2.127	1.173	5.599	5.024	TAG	4.336	2.651	5.580	9.178	1.250	2.522

TAA stop codon environment												
	-5	-4	-3	-2	-1	TAA	+4	+5	+6	+7	+8	+9
A	0.368	0.296	0.387	0.452	0.361	TAA	0.297	0.316	0.368	0.355	0.297	0.393
G	0.161	0.226	0.232	0.097	0.142	TAA	0.187	0.136	0.174	0.122	0.161	0.142
C	0.200	0.239	0.129	0.155	0.181	TAA	0.129	0.200	0.148	0.168	0.271	0.155
T	0.271	0.239	0.252	0.296	0.316	TAA	0.387	0.348	0.310	0.355	0.271	0.310
χ^2	2.358	3.484	6.237	17.687	4.314	TAA	4.559	2.173	1.590	3.310	9.646	3.552

TGA stop codon environment												
	-5	-4	-3	-2	-1	TGA	+4	+5	+6	+7	+8	+9
A	0.348	0.304	0.402	0.424	0.261	TGA	0.347	0.315	0.304	0.391	0.315	0.272
G	0.174	0.239	0.239	0.087	0.163	TGA	0.185	0.196	0.283	0.196	0.174	0.206
C	0.185	0.120	0.152	0.196	0.163	TGA	0.109	0.109	0.163	0.196	0.152	0.185
T	0.293	0.337	0.207	0.293	0.413	TGA	0.359	0.380	0.250	0.217	0.359	0.337
χ^2	0.626	4.244	4.900	9.008	7.980	TGA	2.966	3.641	7.964	4.773	0.720	1.494

The position relative to start or stop codon is indicated at the top of the column. The numbers in the columns give the relative frequency of each base at each position. χ^2 tests were performed with three degrees of freedom (threshold for an α risk of 5% is 7.815 and for an α risk of 1% is 11.345). Expected frequencies used in χ^2 tests are A = 0.32, T = 0.32, G = 0.17 and C = 0.17 in non-coding regions, A = 0.32, G = 0.20, C = 0.19 and T = 0.28 in coding regions. Tabulation performed on 318 ORFs >150 codons.

at positions -3, -2, -1, +4, +5 and +7. We then addressed the question of the possible existence of a consensus sequence in the environment of the stop codons. Not surprisingly, TAA is the more frequently used stop codon: 155 ORFs longer than 150 codons have it, while 92 have

TGA and 71 TAG. When the nucleotide environment between positions -5 and +9 (position +1 being defined by the T of the stop signal) was tabulated, we observed the frequencies reported in Table III. It appears that, in the case of TAA, there is a bias at position -2, which is

more frequently than expected occupied by A and less frequently by G, and at position +8, where C is increased. In the case of TAG, at position -2 the frequency of C is depressed, while this nucleotide is nearly always absent from position +7. Finally, in the case of TGA, the distribution deviates from randomness at three positions, -2, -1 and +6.

Small ORFs (< 100 codons)

The choice of a minimal length of 99 sense codons between the first ATG and the stop signal, which dates back to 1979 (Galibert *et al.*, 1979), probably owes more to the widely used decimal numbering system than to proper insight into biological mechanisms. However, as mentioned above, this size is warranted in the case of yeast (Dujon *et al.*, 1994). In simulation experiments in which chromosome length and nucleotide composition was varied, the chances that ORFs longer than 150 codons will exist and still not correspond to a real gene are negligible. Conversely, the chances that ORFs in the range 100–149 codons will have no biological significance increase in proportion to decreasing size. However, a size of 100 codons is no impassable limit and obviously some ORFs smaller than 100 codons correspond to genes and, for that matter, quite a few proteins shorter than 99 amino acids may not be accounted for by post-translational processing. An example is provided by the small proteolipids PMP1 and PMP2 (40 and 43 amino acids), on chromosomes III and V, respectively (Navarre *et al.*, 1992; 1994). Analysis of the chromosome X sequence has revealed 344 small ORFs 50–98 sense codons in size. Comparison of the deduced proteins with database entries shows that one of these, J0526 (106425–106706), corresponds to the gene encoding subunit VIII of ubiquinol-cytochrome *c* reductase (Hemrika *et al.*, 1993). It is a 94-amino acid protein whose coding gene has been hitherto overlooked. Another instance is YKR057w, which encodes a ribosomal protein of 87 amino acids. Some small ORFs, such as J1567 (479710–479952), J1564 (477910–478074) and J15591 (474126–474368) have perfect or nearly perfect matches with Ty retrotransposon proteins of longer size. These small ORFs most probably result from frameshift mutations, a rather common occurrence in these retrotransposons. Finally, significant similarity is observed between some small ORFs located in the subtelomeric region, such as J0210 (9452–9852), and similar elements located on other chromosomes (K-B110 on chromosome XI or I.A75 on chromosome IX). The other small ORFs, displaying no significant homology with database entries, cannot simply be discarded, since some probably correspond to real genes. Examples in point are J0523 (105893–106060), J1153 (337859–338143), J2123 (676661–676924) and J1425 (448166–448444), all with CAIs >0.2. Clearly, a screening programme taking into account parameters such as the ATG and stop codon environment and the CAI must be developed to approach the question of their existence as genes.

Sequence duplications

We have analysed the nucleotide sequence of chromosome X for the occurrence of sequences demonstrating high similarity to other genes of chromosome X (intrachromosomal duplications) and to genes in other yeast chromo-

somes (interchromosomal duplications), both at the nucleotide and the amino acid level (Table IV). Some of the duplicated ORFs have been functionally characterized. These results confirm earlier observations on chromosomes XI (Dujon *et al.*, 1994) and II (Feldmann *et al.*, 1994) of the high level of internal genetic redundancy in the yeast genome. Moreover, in addition to duplication of individual genes, duplication of syntenic segments has also occurred, syntenic in the present context of intraspecies duplications meaning that two or more genes situated closely on the same chromosome have their homologous loci also located close together, with the same respective orientation, on the other chromosome. As a rule, the physical distance and the nucleotide sequence between two ORFs on the same syntenic segment are not conserved. However, some degree of intergenic sequence conservation can be observed in a few cases, as exemplified in Figure 4.

tRNAs and transposons

Twelve tRNA genes are found on each strand (Figure 5), a density somewhat higher than that observed in the previously sequenced yeast chromosomes. The 24 tRNAs can transfer 13 amino acids in all and include four tRNA^{Asp}, all identical with the same GTC anticodon; four tRNA^{Arg}, two identical with TCT, one with ACG and one with CCT, the last two with minor sequence differences. Of the three tRNA^{Met}, two are identical while the third exhibits slight differences. The two tRNA^{Tyr} have an identical sequence and include the same GTA anticodon.

Upon folding, all the predicted tRNAs fit in readily with the clover-leaf model, regarding stem length as well as loop size. All the canonical bases are observed in all cases but one. The exception is tRNA^{Met} at position 517571, which exhibits an A, instead of T as in the canonical GTYC sequence. Careful checking of the sequence has shown that this ATC sequence does not result from sequencing errors. However, a cloning artefact at some point in the construction of the cosmid library cannot be ruled out at this stage.

While the clover-leaf model is basically respected, 46 non-canonical or unpaired bases are observable in the stems of this two-dimensional configuration. Thirty-nine correspond to a GT base pairing, three to TT and CA and one to GG. An example of such tRNA folding is presented in Figure 6. These observations cannot be ascribed to sequencing or cloning incidents, since they have been observed by different investigators all working on different cosmids. Furthermore, the reality of such pairings has been established by direct RNA sequencing on mature tRNA and by mutagenesis experiments (Pütz *et al.*, 1993). However, it is also true that in the case of plant mitochondrial tRNAs, some (but not all) mismatched base pairs are so edited as to generate a Watson–Crick pair in the mature tRNA (Maréchal-Drouard *et al.*, 1993). While this phenomenon is not yet documented in nuclear yeast tRNA, the possibility of a similar editing process, whereby some of the 46 mispairings mentioned above would be converted into conventional Watson–Crick pairs, cannot be dismissed without additional sequence data or structural studies at the tRNA level. An alternative hypothesis is that some of the predicted tRNAs actually correspond to inactive pseudogenes.

Four of the tRNA genes encountered in chromosome

Table IV. Related genes from chromosome X

Gene/ORF on chromosome X	Related gene/ORF on other chromosome ^a	Functional description ^b	aa identity % ^c	nt identity % ^d
YJL223c	PAU1 (5)	PAU1 protein	96.7 (1-120)/120	96.7 (1-360)/360
YJL210w	LGT3 hexose transport protein	97.9 (1-567)/567	98.4 (883-1701)/1701	
YJL200c	ACO1 (12)	similar to acornitin hydratase	55.3 (35-782)/782	50.8 (6-2278)/2367
YJL198w	YCR037c (3)	probable transport protein	65.0 (39-879)/881	68.1 (684-2387)/2643
YJL196c	YCR034w (3) [*]	similar to sterol isomerase SUR4	58.4 (16-310)/310	60.3 (70-891)/930
YJL191w (CRY2)	CRY1 (3)	ribosomal protein S14eB	96.3 (3-138)/138	92.0 (8-414)/414
YJL190c (RPS24)	L8039.6 (12)	ribosomal protein S15ae	99.2 (1-130)/130	89.1 (1-390)/390
YJL164c (SRA3)	TPK3 (11)	cAMP-dependent protein kinase	84.5 (69-397)/397	73.0 (255-486)/1191
YJL139c (YUR1)	KTR2 (11)	YUR1 protein	66.3 (37-424)/426	64.3 (164-1442)/1284
YJL138c (TIF2)	TIF1 (11)	translation initiation factor eIF-2	100 (1-395)/395	99.3 (1-1185)/1185
YJL133w (MRS3)	MRS4 (11)	mitochondrial splicing protein	76.2 (23-312)/314	70.5 (119-875)/942
YJL099w (CSD3)	YKR027w (11)	CSD3 protein	42.3 (1-844)/1058	37.3 (1759-2238)/2238
YJL098w	YKR028w (11)	unknown	45.8 (1-844)/1058	60.0 (164-1442)/1284
YJL084c	YKR021w (11)	unknown	37.6 (4-932)/1046	46.4 (7-1946)/3138
YJL083w	YKR019c (11)	unknown	26.7 (38-604)/604	64.6 (1265-1601)/1812
YJL082w	YKR018c (11)	unknown	66.0 (1-730)/731	53.7 (233-1986)/1986
YJL079c	YKR013w (11)	unknown	47.5 (1-299)/299	61.4 (415-789)/897
YJL078c	YKR013w (11)	unknown	67.3 (15-161)/881	39.0 (1295-1711)/2643
YJL076w	YKR010c (11)	unknown	16.1 (1-772)/1189	33.7 (2103-3317)/3567
YJL045w	SDH1 (11)	succinate dehydrogenase flavoprotein	83.5 (1-634)/634	78.6 (620-1766)/1902
YJL034w (KAR2)	SSA1 (1)	nuclear fusion protein KAR2 precursor	63.5 (50-663)/682	67.0 (156-1962)/2046
YJL034w (SSC1)	YEL030w (5)	heat shock protein	82.6 (17-642)/654	75.8 (205-1889)/1962
YJR047c (ANB1)	YEL034w (5)	translation initiation factor	90.4 (2-157)/157	91.4 (1-465)/471
YJR048w (CYC1)	YEL039c (5)	cytochrome isoform I	85.8 (2-107)/109	81.9 (113-323)/327
YJR049c (UTR1)	YEL041w (5)	UTR1 protein	57.0 (104-509)/530	63.8 (419-1392)/1590
YJR051w (OSM1)	YEL047c (5)	involved in osmotic regulation	63.5 (36-499)/501	63.7 (218-1469)/1503
YJR066w (TOR1)	TOR2 (11)	phosphatidylinositol kinase	68.0 (62-2470)/2470	67.2 (2786-7410)/7410
YJR103w (URA8)	URA7 (2)	CTP synthase	79.0 (1-562)/564	71.7 (146-1631)/1692
YJR155w	N0300 (14)	similar to aryl-alcohol dehydrogenase	89.9 (1-288)/288	87.7 (1-389)/864
YJR156c	N0295 (14)	similar to thiamine-repressed nmi-1	98.8 (1-340)/340	98.4 (568-1011)/1020
YJL221c	YJL216c	similar to α -glucosidase MAL35 (S46183)	66.3 (11-587)/589	62.8 (199-1767)/1767
YJL219w	YJL214w	similar to hexose transport protein LGT3	65.2 (33-567)/567	66.3 (226-1685)/1701
YJL079c	YJL078c	unknown	66.7 (152-298)/299	66.2 (551-861)/897
YJL052w (TDH1)	YJR009c (TDH2)	glyceraldehyde-3-phosphate dehydrogenase	65.0 (1-331)/331	92.4 (1-996)/996
YJL038c	YJL037w	unknown	36.3 (5-218)/219	34.0 (295-640)/657

^aWhere known, chromosomal location is indicated in parenthesis.

^bFunction of genes on chromosome X, when available, or else function of their homologues on other chromosomes.

^cNumbers indicate % of aa identity, boundaries of aa comparison (in brackets) and size of the ORF on chromosome X (number after dash).

^dSame as above, but in nt.




Fig. 4. Physical comparison of the location of genes and syntenic segments on chromosome X with that of their counterparts on other chromosomes. The precise position of the genes was deduced from the present sequence and re-drawn to scale (coordinates are in kb). Elements above and below the scale belong to the Watson and the Crick strands, respectively. Shaded boxes represent the ORFs with a counterpart on the other chromosome. On the whole, physical distance (and the structures located therein) between any two ORFs in the same syntenic segment is not respected on chromosomes other than X. Exceptions are the consecutive ORFs YJL099w (CSD3) and YJL098w on chromosome X and their homologues YKR027w and YKR028w on chromosome XI, the consecutive ORFs YJL083w and YJL082w on chromosome X and their homologues YKR019c and YKR018c.




Fig. 5. Position of tRNA genes, Ty sequences and LTRs on chromosome X. The positions were drawn to scale relative to the complete sequence. Elements on the Watson and Crick strands are displayed on the right- and left-hand side, respectively. Only the 5' coordinate is given.




Fig. 6. A clover-leaf structure of yeast tRNA^{Met} on chromosome X (422 624–422 696). All canonical bases are indicated by asterisks. Mismatched base pairs in the stems are boxed. The shadowed nucleotides are the anticodon.

X display an intron 3' to the anticodon sequence, as previously observed. These include two tRNA^{Tyr} with an intron of 14 nt, one of the two tRNA^{Leu} with a 19-nt intron and the unique tRNA^{Trp} with an intron of ~29 nt. Its exact size is difficult to assess because base pairing is possible between several short sequences in the anticodon stem, creating an extra arm of variable length.

The entire chromosome X sequence was scanned in parallel for the presence of complete Ty elements or solo remnants or LTR thereof. As shown in Figure 5, several of these have been found. One complete Ty4 is present at position 197243–203468 and two complete Ty1 at position 472150–483659. The two elements are in tandem and share a central δ element. In addition, several solo LTRs are observed. As reported, with the exception of Ty1 these elements are located in the vicinity of tRNA sequences. However, this association seems to be rather loose and, besides, it involves partners located on either strand relative to one another.

Comparison of the physical and genetic maps of the chromosome X

The genetic map of chromosome X includes 60 genes or markers, of which 48 were mapped in a linear array and 12 remained unmapped (Mortimer *et al.*, 1995). Figure 7 shows a comparison of this map with the physical map deduced from the complete nucleotide sequence. Contrary to what has been reported for chromosome XI (Dujon *et al.*, 1994), no gross translocation or inversion was observed here. On the whole, the intergenic distance on the genetic map is roughly proportional to the physical distance, indicative of a relatively uniform recombination frequency over chromosome X. However, closer examination reveals some interesting discrepancies. First, genetic mapping has assigned the previously sequenced *CYR1* gene (alias *CDC35*, *HSR1*, *SRA4* and *TSM0185*), encoding adenylyl cyclase, to a site indistinguishable from that of *sui2*. This assignment is clearly incorrect, as the sequence data shows that this gene is in fact located on the left arm of the chromosome, close to the centromere. Second, marked differences are observed in map distances, the ratio between genetic and physical map distances ranging from 0.02 cM per kb for the *TDH2/met3* marker pair, to 0.84 and 4.74 cM per kb for the *met3/ilv3* and *ilv3/ess1* pairs, respectively. The relatively high frequency of recombination observed in these latter intervals strongly suggests the existence of preferred sites for the initiation of meiotic recombination, similar to those found in the *arg4* region on chromosome VIII (Nicolas *et al.*, 1989; Sun *et al.*, 1989) and the *MAT/thr4* region on chromosome III (Jacquet *et al.*, 1991). It is interesting to note that these intervals of high recombination frequencies in chromosome X appear to coincide with the sharp peak in the G+C content in the right arm of the chromosome (Figure 2).

In all, 31 of the mapped and one, tRNA^{Ser}, of the unmapped could be unambiguously assigned to an ORF or a tRNA gene on the basis of sequence comparison. A total of 28 loci cannot at present be attributed to specific ORFs on the physical map of chromosome X.

Discussion

The various elements of the chromosome X sequence referred to above are depicted in Figure 8. The present




Fig. 7. Comparison of the genetic and physical maps of yeast chromosome X. The genetic map is re-drawn from Mortimer (Mortimer *et al.*, 1995). The unmapped genes or markers are listed on the right. The physical map deduced from this work has been drawn to scale. The circle indicates the position of the centromere. Genes or markers for which no corresponding ORF has been identified on the physical map are indicated by an asterisk.

report brings the number of completely sequenced chromosomes from the yeast *S.cerevisiae* to nine, chromosome X ranking second in this series by virtue of its size. Thus, nearly 40% of the *S.cerevisiae* genome sequence is now accessible to analysis, availability of the whole sequence being anticipated for 1997. The sequence of chromosome X has been established in S288C, a *S.cerevisiae* strain chosen by all members of the European Union sequencing consortium led by André Goffeau. While the study of this sequence reveals no features that are specific for chromosome X, it corroborates several observations made with the previously sequenced chromosomes.

Taking into account only those ORFs whose characteristics, such as size, CAI and disposition leave no doubt as to their existence as real genes, a minimal density of one gene per 2000 nt can be estimated. All these genes are regularly spaced along the chromosome, with no predilection for either strand. Following translation and comparison of the deduced amino acid sequence with database entries, the products of these ORFs can be categorized as follows: (i) 102 proteins previously identified in *S.cerevisiae* and encoded by genes already assigned to chromosome X; (ii) 16 proteins with strong similarity,


or even near identity, to known *S.cerevisiae* proteins, but whose coding gene has not previously been shown to reside on chromosome X; (iii) 22 proteins with a FastA score much greater than 200—equal to at least half the self-score, i.e. the score obtained when the protein is compared with itself. Such high scores can be considered as warranting a realistic hypothesis regarding the function of ORFs in this category; (iv) 35 proteins with a FastA score >200, though lower than half the self-score. A function can also be envisaged in this case, but with more caution; (v) 92 proteins with no significant FastA score but displaying a particular motif signature; (vi) 112 proteins with no match at all in database entries. This last category remains numerically important, since it includes nearly 30% of the ORFs, a proportion that fully vindicates the systematic sequencing approach of the *S.cerevisiae* genome launched in 1989.

Regarding ORFs in categories (iii) and (iv) above, for which a function can be hypothesized, several of the proteins discovered in chromosome X are worth mentioning. For instance, three new genes encoding different subunits of the cytosolic chaperone complex (*CCT5*, *CCT7* and *CCT8*) have been discovered on chromosome X in addition to *CCT3*. This brings the number of fully sequenced *CCT* genes in *S.cerevisiae* to eight. Together with the versatility of yeast versus mouse genetics, availability of these sequence data will undoubtedly promote fine molecular analysis of this important chaperone system. Another remark concerns the discovery of a Cl⁻ channel gene (Huang *et al.*, 1994e) on chromosome X. In this respect, it is both surprising and remarkable that systematic sequencing was required to detect the first Cl⁻ channel ever described in a species as thoroughly studied as *S.cerevisiae*. Here again, availability of the gene and of disruption mutants thereof will permit identification by complementation homologous genes in other species of interest, in particular in plants.

Chromosome X stands out because of the number of tRNA genes (24) it accommodates, capable of transferring 13 different amino acids. However, what is even more remarkable and has so far escaped notice is that folding of these tRNAs according to the clover-leaf model reveals quite a few mismatches in the several stems. This is suggestive of an editing process aiming at correcting some of these mismatches, as reported for various tRNAs from plants (Maréchal-Drouard, 1993). Of course, validation or dismissal of this hypothesis must await analysis at the RNA level.

Duplicated genes are found in chromosome X, as in other *S.cerevisiae* chromosomes. These include both intra- and interchromosomal duplications. Furthermore, actual syntenic regions can be recognized in the latter case. The implications are 2-fold, pertaining (i) to the study of the evolution of the yeast genome and (ii) to function analysis, as it is known that disruption of a single gene frequently does not result in any phenotypic alteration. By the same token, a clue to the function of a gene might in some instances be provided by disruption of all the genes belonging to a given family.

To conclude, it must be stressed that this brief account of the sequence analysis of chromosome X cannot cover all the information embedded in the nucleotide sequence



and that many biological analyses will be needed to exploit this mine of information in the years to come.

Materials and methods

Chromosome X DNA

Total yeast DNA was obtained from FY1679, a diploid strain issued from the cross between strains FY23 (*MATa, ura3-52, trp1Δ63, leu2Δ1, GAL2*) and FY73 (*MATα, ura3-52, his 3Δ200, GAL2*). FY23 and FY73 are derived from strain S288C and are isogenic with it except for the

markers indicated (Winston *et al.*, 1995). The construction of an ordered cosmid library and of an *EcoRI* restriction map have been previously published (Huang *et al.*, 1994a). Overlapping cosmids covering the chromosome X contig were distributed within a consortium of 15 laboratories. The telomeres and subtelomeric regions were cloned in vector pEI.61, as described by Louis and Borts (1995).

Determination, assembly and analysis of the sequence

Sequencing strategies and methods varied among the 15 collaborating laboratories (Table V). Sequence assembly in the single contracting laboratories was performed by a variety of software program packages.




Fig. 8. Chromosome X map deduced from the complete sequence. The chromosome and its constitutive elements are drawn to scale. The top bar represents the Watson strand oriented 5' to 3' from left to right, the bottom bar the Crick strand. The conserved elements of the centromere are designated as CDE I, II and III. ORFs on the left and right arm are designated by the letters L and R, respectively, before their number (numbering is in increasing order from the centromere). Full designations, in accordance with the official ORF nomenclature, are obtained by adding again the letters Y (for yeast) and J (for chromosome X) at the beginning, and w (Watson) or c (Crick) at the end.

The telomeres were cloned in Oxford. The left telomere was sequenced in one of 15 laboratories. The right telomere and the PCR fragment filling the gap were sequenced in Berlin. Completed contigs submitted to MIPS were stored in a data library and assembled using the GCG software package 7.2 for the VAX (Devreux *et al.*, 1984). The nature

and position of genetic elements have been deduced from the sequence using the following principles: (i) all possible intron splice site/branch-point pairs were detected using specially defined patterns (Fondrat *et al.*, 1994); (ii) ORFs occurring in all possible frames were listed. ORFs containing at least 99 contiguous sense codons following an ATG and

Table V. Methods used by each of the collaborating laboratories

Whole cosmid Shotgun	Restricted fragments		
	Shotgun	TN1000	Nested deletions
Louvain (M) Heidelberg (M) Konstanz (M) Paris (A) Gif (A) Rennes (A)	Gembloux (M) Amsterdam (A)	Darmstadt (M) Frankfurt (A)	München (A) Copenhagen (A) Düsseldorf (A) Ghent (A) Herakleion (M)

M, manual methods; A, automated methods.

those containing 50–98 codons were retained for further analysis, in both cases provided they were not entirely contained within a longer ORF on either DNA strand. Searches for similarity of the deduced protein sequences to entries in the databanks were performed by FastA (Pearson and Lipman, 1988) in the Protein Sequence Database of PIR International (release 44) and other databases. Protein signatures were detected using the PROSITE dictionary (release 11.1) (Bairoch, 1989). ORFs were assigned probable functions when the alignments from FastA searches showed significant similarity and/or protein signatures were apparent, whereas FastA scores <200 were considered insufficient to confidently assign function. The complete sequence was also searched for tRNA genes ('trnscan') (Fichant and Burks, 1991), centromere and telomere consensus elements and for δ , σ or τ elements by comparison with a data set of such elements previously characterized in yeast. Compositional analyses of the chromosome were performed using the XII program package (C.Marck, unpublished results). For calculations of CAI and GC content of ORFs, the algorithm CODONS (Lloyd and Sharp, 1992) was used.

Sequence verifications and quality controls

All sequences submitted by collaborating laboratories to the Martinsried Institute for Protein Sequences (MIPS) data library were subjected to quality controls. The procedure was comprised of three major steps. First, the strategy of each contractor was checked by the coordinator to pinpoint possible weak points and request the sequencers to review their electrophoretograms to assess the quality of their reads in these less documented regions. Second, once cosmid sequences had been entered in the database, the match between the overlaps was held to provide an assessment of the respective quality of the neighboring partial sequences. Third, each of the cosmids that had been distributed to the contractors for sequencing was shotgunned, size-selected to ~300–500 bp and cloned in plasmid vector, the size of the inserts ensuring that sequencing with the universal forward and reverse primers would provide a 300–400 double-stranded sequence. The subclones from each cosmid were sent with coded names to a different sequencer. The double-stranded part of each sequence was then sent to MIPS and compared with the initial sequence. The number of verification sequences per cosmid clone (averaging 15–30) varied according to the quality of the initial sequencing as deduced from alignment within the overlaps. Any discrepancy detected between overlapping partial sequences or between the sequence initially submitted and the verification sequence was addressed as follows. A stretch of 20 bp including the discrepancy, but not centering on it, was pointed out to each party for reviewing and re-submission to MIPS, whether modified or not. This procedure was sufficient to remove most discrepancies, as one party usually provided a revised sequence matching the other's. Resistant cases were dealt with by requesting both parties to send the electrophoretograms corresponding to the conflicting sequences to the coordinator, who made a decision and requested resequencing if necessary.

The sequence data reported are available through <http://mips.biochem.mpg.de/yeast>

Acknowledgements

We wish to thank B.Dujon for fruitful discussion and for help with the gene density and G+C composition plots, and G.Le Provost for secretarial assistance. The laboratory consortium operating under contracts with the European Commission was initiated and organized by A.Goffeau. This study is part of the second phase of the European Yeast Genome Sequencing Project carried out under the administrative coordination of

A.Vassarotti (DG-XII) and the Université Catholique de Louvain, and under the scientific responsibility of F.Galibert as DNA coordinator. This work was supported by the European Commission under the BRIDGE and Biotech programmes, the Groupe de Recherche et d'Etudes sur le Génome (GREG) and the Centre National de la Recherche Scientifique (CNRS) (FR), the Wellcome Trust (UK), the Région Wallonne and the Fonds National de la Recherche Scientifique (BE), the Bundesminister für Forschung und Technologie (DE) and the Ministry of Industry and Technology (GR).

References

- Bairoch, A. (1989) EMBL Biocomputing Technical Document 4. EMBL, Heidelberg, Germany.
- Barrell, B.G. et al. (1994) Sequence of *S. cerevisiae* chromosome IX. <http://www.sanger.ac.uk/~yeast/pub/svw/seqnecing.html>
- Bussey, H. et al. (1995) The nucleotide sequence of chromosome I from *Saccharomyces cerevisiae*. *Proc. Natl Acad. Sci. USA*, **92**, 3809–3813.
- Delehotde, A., Goguel, V., Becam, A.M., Creusot, F., Perea, J., Banroques, J. and Jacq, C. (1989) Site-specific DNA endonuclease and RNA maturase activities of two homologous intron-encoded proteins from yeast mitochondria. *Cell*, **56**, 431–441.
- Deverenz, J., Haeberli, P. and Smithies, O. (1984) A comprehensive set of sequence analysis programs for the VAX. *Nucleic Acids Res.*, **12**, 387–395.
- Dietrich, F.S. et al. (1994) Sequence of *S. cerevisiae* chromosome V. <http://speedy.mips.biochem.mpg.de/mips/yeast/chr5>
- Dujon, B. et al. (1994) Complete DNA sequence of yeast chromosome XI. *Nature*, **369**, 371–378.
- Feldmann, H. et al. (1994) Complete DNA sequence of yeast chromosome II. *EMBO J.*, **13**, 5795–5809.
- Fichant, G.A. and Burks, C. (1991) Identifying potential tRNA genes in genomic DNA sequences. *J. Mol. Biol.*, **220**, 659–671.
- Fondrat, C. and Kalogeropoulos, A. (1994) Approaching the function of new genes by detection of their potential upstream activation sequences in *Saccharomyces cerevisiae*: application to chromosome III. *Curr. Genet.*, **25**, 396–406.
- Galibert, F., Mandart, E., Fitoussi, F., Tiollais, P. and Charnay, P. (1979) Nucleotide sequence of the hepatitis B virus genome (subtype ayw) cloned in *E. coli*. *Nature*, **281**, 646–650.
- Hemrika, W., Berden, J.A. and Grivell, L.A. (1993) A region of the C-terminal part of the 11-kDa subunit of ubiquinol-cytochrome-c oxidoreductase of the yeast *Saccharomyces cerevisiae* contributes to the structure of the Q(ont) reaction domain. *Eur. J. Biochem.*, **215**, 601–609.
- Hieter, P., Pridmore, D., Hegemann, J.H., Thomas, M., Davis, R.W. and Philippsen, P. (1985) Functional selection and analysis of yeast centromeric DNA. *Cell*, **42**, 913–921.
- Hinnebusch, A.G. and Liebman, S.W. (1991) Protein synthesis and translational control in *Saccharomyces cerevisiae*. In Broach, J.R. et al. (eds), *The Molecular Biology of the Yeast Saccharomyces*. Cold Spring Harbor Laboratory Press, Plainview, NY, pp. 627–735.
- Huang, M.E., Chnat, J.C., Thierry, A., Dujon, B. and Galibert, F. (1994a) Construction of a cosmid contig and of an EcoRI restriction map of yeast chromosome X. *DNA Sequence*, **4**, 293–300.
- Huang, M.E., Manus, V., Chuat, J.C. and Galibert, F. (1994b) Revised nucleotide sequence of the COR region of yeast *S. cerevisiae* chromosome X. *Yeast*, **10**, 811–818.

- Huang, M.E., Chuat, J.C. and Galibert, F. (1994c) A voltage-gated chloride channel in the yeast *Saccharomyces cerevisiae*. *J. Mol. Biol.*, **242**, 595–598.
- Huang, M.E., Chuat, J.C. and Galibert, F. (1995) Analysis of a 42.5 kb DNA sequence of chromosome X reveals three tRNA genes and 14 new open reading frames including a gene most probably belonging to the family of ubiquitin-protein ligase. *Yeast*, **11**, 775–781.
- Jacquet, M., Buhler, J.-M., Iborra, F., Francinques-Gaillard, M.C. and Soustelle, C. (1991) The *MAT* locus revisited within a 9.8 kb fragment of chromosome III containing *BUD5* and two new open reading frames. *Yeast*, **7**, 881–888.
- Johnston, M. et al. (1994) Complete nucleotide sequence of *Saccharomyces cerevisiae* chromosome VIII. *Science*, **265**, 2077–2082.
- Ketchum, K.A., Joiner, W.J., Sellers, A.J., Kaczmarek, L.K. and Goldstein, S.A.N. (1995) A new family of outwardly rectifying potassium channel proteins with two pore domains in tandem. *Nature*, **376**, 690–695.
- Kozak, M. (1987) An analysis of 5′-noncoding sequences from 699 vertebrate messenger RNAs. *Nucleic Acids Res.*, **15**, 8125–8148.
- Lloyd, A.T. and Sharp, P.M. (1992) CODONS: A microcomputer program for codon usage analysis. *J. Hered.*, **83**, 239–240.
- Louis, E.J. and Borts, R.H. (1995) A complete set of marked telomeres in *Saccharomyces cerevisiae* for physical mapping and cloning. *Genetics*, **139**, 125–136.
- Louis, E.J. and Haber, J.E. (1991) Evolutionarily recent transfer of a group I mitochondrial intron to telomere regions in *Saccharomyces cerevisiae*. *Curr. Genet.*, **20**, 411–415.
- Louis, E.J., Naumova, E.S., Lee, A., Naumov, G. and Haber, J.E. (1994) The chromosome end in yeast: its mosaic nature and influence on recombinational dynamics. *Genetics*, **136**, 789–802.
- Maréchal-Drouard, L., Ramamonjisoa, D., Cosset, A., Weil, J.H. and Dietrich, A. (1993) Editing correct mispairing in the acceptor stem of bean and potato mitochondrial phenylalanine transfer RNAs. *Nucleic Acids Res.*, **21**, 4909–4914.
- Miosga, T., Witzel, A. and Zimmermann, F.K. (1994a) Sequence and function analysis of a 9.46 kb fragment of *Saccharomyces cerevisiae* chromosome X. *Yeast*, **10**, 965–973.
- Miosga, T., Boles, E., Schaaff-Gerstenschläger, I., Schmitt, S. and Zimmermann, F.K. (1994b) Sequence and function analysis of a 9.74 kb fragment of *Saccharomyces cerevisiae* chromosome X including the BCK1 gene. *Yeast*, **10**, 1481–1488.
- Miosga, T., Schaaff-Gerstenschläger, I., Chalwatzis, N., Baur, A., Boles, E., Faarnier, C., Schmitt, S., Velten, C., Wilhelm, N. and Zimmermann, F.K. (1995) Sequence analysis of a 33.1 kb fragment from the left arm of *Saccharomyces cerevisiae* chromosome X, including putative proteins with leucine zippers, a fungal Zn(II)-Cys6 binuclear cluster domain and a putative alpha2-SCB-binding site. *Yeast*, **11**, 681–689.
- Mortimer, R.K., Cherry, J.M., Dietrich, F.S., Riles, L., Olson, M.S. and Botstein, D. (1995) Genetic map of *Saccharomyces cerevisiae*. Edition 17. <http://genome-www.stanford.edu/sacchdb/edition12.html>
- Murakami, Y. et al. (1995) Analysis of the nucleotide sequence of chromosome VI from *Saccharomyces cerevisiae*. *Nature Genet.*, **10**, 261–268.
- Navarre, C., Ghislin, M., Leterme, S., Ferroud, C., Dufour, J.P. and Goffeau, A. (1992) Purification and complete sequence of a small proteolipid associated with the plasma membrane H(+)-ATPase of *Saccharomyces cerevisiae*. *J. Biol. Chem.*, **267**, 6425–6428.
- Navarre, C., Catty, P., Leterme, S., Dietrich, F. and Goffeau, A. (1994) Two distinct genes encode small isoproteolipids affecting plasma membrane H(+)-ATPase activity of *Saccharomyces cerevisiae*. *J. Biol. Chem.*, **269**, 21262–21268.
- Nicolas, A., Treco, D., Schultes, N.P. and Szostak, J.W. (1989) An initiation site for meiotic gene conversion in the yeast *Saccharomyces cerevisiae*. *Nature*, **333**, 87–90.
- Oliver, S. et al. (1992) The complete DNA sequence of yeast chromosome III. *Nature*, **357**, 38–46.
- Pearson, W.R. and Lipman, D.J. (1988) Improved tools for biological sequence comparison. *Proc. Natl Acad. Sci. USA*, **85**, 2444–2448.
- Pryde, F.E., Huckle, T.C. and Louis, E.J. (1995) Sequence analysis of the right end of chromosome XV in *Saccharomyces cerevisiae*: An insight into the structural and functional significance of sub-telomeric repeat sequences. *Yeast*, **11**, 371–382.
- Purnelle, B., Coster, F. and Goffeau, A. (1994) The sequence of a 36 kb segment on the left arm of yeast chromosome X identifies 24 open reading frames including *NUC1*, *PRP21(SPP91)*, *CDC6*, *CRY2*, the gene for S24, a homologue to the aconitase gene *ACO1* and two homologues to chromosome III genes. *Yeast*, **10**, 1235–1249.
- Pütz, J., Puglisi, J.D., Florentz, C. and Giegé, R. (1993) Addictive, cooperative and anti-cooperative effects between identity nucleotides of a tRNA. *EMBO J.*, **12**, 2949–2951.
- Rasmussen, S.W. (1995) A region of yeast chromosome X includes the *SME1*, *MEF2*, *GSH1* and *CSD3* genes, a TCP-1 related gene, an open reading frame similar to the *DAL80* gene, and a tRNA^{Met}. *Yeast*, **11**, 873–883.
- Sharp, P.M. and Li, W.-H. (1987) The codon adaptation index — a measure of directional synonymous codon usage bias, and its potential applications. *Nucleic Acids Res.*, **15**, 1281–1295.
- Sun, H., Treco, D., Scultes, N.P. and Szostak, J.W. (1989) Double stranded breaks at an initiation site for meiotic gene conversions. *Nature*, **333**, 87–90.
- Vandenbol, M., Durand, P., Bolle, P.-A., Dion, C., Portetelle, D. and Hilger, F. (1994) Sequence analysis of a 40.2 kb DNA fragment located near the left telomere of yeast chromosome X. *Yeast*, **10**, 1657–1662.
- Vandenbol, M., Durand, P., Dion, C., Portetelle, D. and Hilger, F. (1995) Sequence of a 17.1 kb DNA fragment from chromosome X of *Saccharomyces cerevisiae* includes the mitochondrial ribosomal protein L8. *Yeast*, **11**, 57–60.
- Winston, F., Dollard, C. and Ricuperdo-Hovasse, S.L. (1995) Construction of a set of convenient *Saccharomyces cerevisiae* strains that are isogenic to S288C. *Yeast*, **11**, 53–55.
- Zagulski, M., Babinska, B., Gromadka, R., Migdalski, A., Sulicka, J. and Herbert, C.J. (1995) The sequence of 24.3 kb from chromosome X reveals 5 complete open reading frames all of which correspond to new genes, and a tandem insertion of a Ty1 transposon. *Yeast*, **11**, 1179–1186.

Received on November 3, 1995; revised on January 5, 1996